




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/00</b>		(11) International Publication Number: <b>WO 99/58652</b>
<b>A2</b>		(43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/DK99/00258 (22) International Filing Date: 7 May 1999 (07.05.99) (30) Priority Data: 0627/98                      7 May 1998 (07.05.98)                      DK 60/085,067                  12 May 1998 (12.05.98)                      US (71) Applicant (for all designated States except US): GX BIOSYS-TEMS A/S [DK/DK]; Forskerbyen Symbion, Fruebjergvej 3, DK-2100 Copenhagen Ø (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): GERDES, Kenn [DK/DK]; Majsmarken 17, DK-5260 Odense S (DK). GOTFREDSEN, Marie [DK/DK]; Ørstedsgade 44, 2. th, DK-5000 Odense C (DK). GRØNLUND, Hugo [DK/DK]; Uffesvej 26, DK-5200 Odense V (DK). PEDERSEN, Kim [DK/DK]; Solfaldsvej 18-11, DK-5000 Odense C (DK). KRISTOFFERSEN, Peter [DK/DK]; Bispeengen 15, 1. tv, DK-2000 Frederiksberg (DK). (74) Agent: PLOUGMANN, VINGTOFT & PARTNERS A/S; Sankt Annae Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published Without international search report and to be republished upon receipt of that report. With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.

(54) Title: CYTOTOXIN-BASED BIOLOGICAL CONTAINMENT

## (57) Abstract

Method of conditionally controlling the survivability of a recombinant cell population and of containing such cells to an environment or containing replicons to a host cell is based on the use of proteic killer systems including the *E. coli relBE* locus and similar systems found in Gram-negative and Gram-positive bacteria and *Archae*. Such system are generally based on a cytotoxin polypeptide and an antitoxin or antidote polypeptide that in contrast to the cytotoxin is degradable by proteases. The recombinant cells are useful as vaccines, pollutant degrading organisms or as biological pest control organisms e.g. expressing *B. thuringiensis* crystalline proteins.



09/472,654

BE

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## CYTOTOXIN-BASED BIOLOGICAL CONTAINMENT

## FIELD OF THE INVENTION

5

The present invention relates to the field of biologically containing genetically modified microorganisms in a particular environment and vectors to a particular host cell. Specifically, there is provided recombinant vectors and cells containing a proteic killer system based on the *E. coli* RelE polypeptide and functional equivalents of this cytotoxin and method of containing replicons and cells, respectively to particular host cells or particular environments, respectively.

10

## TECHNICAL BACKGROUND AND PRIOR ART

15

The increasing application of recombinant DNA technology to engineer novel microorganism which are industrially useful have caused concerns in the general public over the potential risks involved. These concerns are primarily related to the potential harm to humans and to undesirable and/or uncontrollable ecological consequences upon de-

20 liberate or unintentional release of such genetically engineered microorganisms (GEMs) into the environment. These concerns have led to the establishment of official guidelines for the safe handling of GEMs in laboratories and production facilities where such organisms are applied. Up till now, such guidelines have primarily been directed to measures of physically containing GEMs in laboratories and production facilities with

25 the aim of reducing the likelihood that workers in such facilities were contaminated, or that the GEMs were to escape from their primary physical environment, such as a fermentation vessel.

30

It is presently being recognized that the level of safety in the handling of GEMs can be increased by combining physical containment measures with biological containment measures to reduce the possibility of the survival of the genetically engineered organisms if they were to escape from their primary environment.

Lately, however, concerns have become increasingly focused on potential risks related to deliberate release of GEMs to the outer environment and to the use of GEMs as live vaccines. In this connection there is a strongly felt need to have biological containment systems which subsequent to the environmental release of the GEMs or their  
5 administration as vaccines to a human or an animal body, effectively kill the released organisms in a controlled manner or which limit the function of the released GEMs to an extent where such GEMs are placed at a significant competitive disadvantage whereby they will eventually be ousted by the natural microflora of the environment to which they are released.

10

The first systems of biological containment were based on the use of "safe" cloning vectors and debilitated host bacteria. As examples, it has been suggested to select vectors which lack transfer functions or which naturally have a very narrow host range. Examples of debilitated host bacteria are *E. coli* mutants having an obligate re-  
15 quirement for exogenous nutrients not present or present in low concentrations outside the primary environment of the GEMs.

Other suggested biological containment systems have been based on mechanisms whereby the vector is restricted to the GEMs e.g. by using a plasmid vector with a  
20 nonsense mutation in a gene, the expression of which is indispensable for plasmid replication or a suppressor mutation in the chromosome, said mutation blocking translational read-through of the message of the gene. A further approach is to maintain the rDNA stably in the host by integrating it into the chromosomes of the GEMs.

25 Recently, an alternative biological containment strategy has been developed in which a recombinant vector is endowed with a gene encoding a cell killing function which gene is under the control of a promoter only being expressed under certain environmental conditions, such as conditions prevailing in an environment outside the primary environment of the GEMs, or when the vector is unintentionally transferred to a secondary  
30 host, or the expression of which is stochastically induced. By using incorporation in a GEM of such a cell killing function and selecting appropriate regulatory sequences, vectors can be constructed which are contained in the primary host cell and/or in a primary physical environment. A cell killing function as defined herein may also be referred to as an active biological containment factor.

If a stochastically induced mechanism of expression regulation is selected for such a biological containment system, a population of GEMs containing the system will, upon release to the outer environment or if used as a live vaccine, be subjected to a random cell killing which will lead to an increase of the doubling time of the host cell population or eventually to the disappearance of the organisms.

The above-mentioned genes encoding cell killing functions are also frequently referred to as "suicide" genes, and biological containment systems based upon the use of such genes, the expression of which are regulated as defined above, are commonly described as conditional lethal systems or "suicide" systems. Up till now, several such cell killing functions have been found in bacterial chromosomes and in prokaryotic plasmids. Examples of chromosomal genes having cell killing functions are the *gef* (Poulsen et al., 1990) and *relF* (Bech et al., 1985) genes from *E. coli* K-12. Examples of plasmid encoded suicide genes are *hok* and *flmA* (Gerdes et al., 1986) genes isolated from plasmids R1 and F, respectively, the *snrB* gene also isolated from plasmid F (Akimoto et al., 1986) and the *pnd* gene isolated from plasmids R16 and R483 (Sakikawa et al., 1989 and Ono et al., 1987). Common features of these genes are that they are transcribed constitutively, regulated at a post-transcriptional level, and that they all encode small toxic proteins of about 50 amino acids and that their translation is controlled by antisense RNA. The application of the *hok* gene in a biological containment system has been disclosed in WO 87/05932.

An alternative biological containment system is disclosed in WO 95/10614 which is based on the use of genes, the expression of which in a cell where the gene is inserted, results in the formation of mature forms of exoenzymes which are hydrolytically active in the cytoplasm of the cell and which can not be transported over the cell membrane. When such enzymes are expressed, the normal function of the cell becomes limited to an extent whereby the competitiveness, and hence the survival, of a population of such cells is reduced.

30

The stable maintenance of low copy-number plasmids in bacteria is secured by a number of plasmid-borne gene systems, one of which is based on killing of plasmid-free cells (also termed post-segregational killing). This regulated killing is based on a toxin-antidote principle, i.e. a two-component system comprising a stable toxin and an un-

stable antidote for the toxin. One such system, which is referred to as a proteic killer gene system is based on protein toxins and protein antidotes (reviewed by Jensen and Gerdes, 1995). The natural function of such systems is to provide stable maintenance of plasmids and it has not been suggested previously to utilize the systems as the basis for confining GEMs to a particular environment.

The *E. coli relB* operon encodes three genes, *relB*, *relE* and *relF* (Bech et al., 1985). It has now been found that *relE* encodes a cytotoxin whose overproduction is lethal to host cells and that the *relB* gene encodes an antitoxin that prevents the lethal action of RelE. When present on a plasmid, the *relBE* operon was able to stabilize the inheritance of a mini-R1 test plasmid. It was also found that *relBE* homologous gene systems are found in a wide variety of Gram-negative and Gram-positive bacteria and in *Archae*.

These results show that the *relBE* genes constitute a new ubiquitously occurring family of gene systems that belongs to the proteic plasmid stabilization systems.

These findings has opened up for an alternative, highly effective and versatile biological containment system as it is described in the following. Importantly, it has been discovered that such a system involves the significant advantage that the frequency of spontaneously occurring mutants of microorganisms that have become resistant to the lethal effect of these cytotoxins is very low. This implies that this biological containment system is very safe.

## SUMMARY OF THE INVENTION

Accordingly, the invention pertains in a first aspect to a method of conditionally controlling the survivability of a recombinant microbial cell population, the method comprising (i) providing in the cells of said population a gene coding for a cytotoxic first kind of polypeptide, the gene is selected from the group consisting of the gene coding for the *E. coli* K-12 RelE polypeptide and a gene coding for a functionally equivalent polypeptide (said genes collectively being designated herein as the *relE* gene family), said gene is expressible in the cells of the population and, operably linked to the gene,

a regulatable regulatory DNA sequence and (ii) cultivating the cell population under conditions where the *relE* gene or the gene coding for a functionally equivalent polypeptide is expressed; the expression leading to an at least partial killing of the cell population.

5

In a further aspect there is provided a method of confining an extrachromosomal replicon to a microbial cell population, the method comprising the steps of

(i) isolating a microbial cell naturally containing a gene belonging to the *relE* gene family coding for a first kind of polypeptide that, when it is expressed in the cell, acts as a toxin for the cell or, if the cell does not naturally contain a gene belonging to the *relE* gene family, introducing such a gene into the cell,

(ii) introducing into the cell the extrachromosomal replicon to be confined, said replicon containing a gene coding for a second kind of polypeptide that, by binding to the first kind of polypeptide, acts as an antitoxin for said first kind of polypeptide,

(iii) cultivating the cell under conditions where the genes coding for the first and the second kind of polypeptides are expressed, whereby a daughter cell that does not receive a copy of the extrachromosomal replicon is killed by the first kind of polypeptide being expressed in the absence of expression of the second kind of polypeptide.

In a still further aspect, the invention relates to a method of post-segregationally stabilizing a plasmid in a microbial host cell population, the method comprising the steps of

(i) recombinationally inserting into the plasmid (a) a gene coding for a first kind of polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide and a functional equivalent thereof, said first kind of polypeptide having a toxin effect on the host cell and (b) a gene coding for a second kind of polypeptide that (1) is capable of acting as an antitoxin for first kind of polypeptide and (2) is capable of being degraded in the host cell at a higher rate than that at which the first kind of polypeptide is degraded,

(ii) cultivating the cell population under conditions where the genes coding for the first kind and second kind of polypeptides are expressed,

whereby a daughter cell that does not receive at least one copy of the plasmid is killed  
5 as a result of the faster degradation of the second kind of polypeptide.

In yet other aspects, the invention provides a recombinant microbial cell comprising a gene coding for a first kind of polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide, a gene coding for a functionally equivalent polypeptide  
10 hereof or a variant or derivative of any such polypeptide, said first kind of polypeptide having a toxic effect on the cell, subject to the limitation that when the cell is *E. coli*, the gene coding for the first kind of polypeptide is not derived from *E. coli*, and a composition comprising such cells.

15 The invention also pertains to several methods of containing cells or replicons including

(1) a method of limiting the survival of a cell population in a first or a second environment, which method comprises

20

(i) transforming the cells of said population with a gene coding for a cytotoxic polypeptide, the gene is selected from the group consisting of the gene coding for the *E. coli* K-12 RelE polypeptide, the gene coding for the plasmid F CcdB polypeptide, the gene coding for the plasmid R1 PemK polypeptide, the gene coding for plasmid RP4  
25 ParE polypeptide, the gene coding for the prophage P1 Doc polypeptide and a gene coding for a functionally equivalent polypeptide for anyone of said polypeptides, said gene is expressible in the cells of the population, and operably linked to the gene, a regulatory DNA sequence being regulatable by an environmental factor and which regulates the expression of said gene, and

30

(ii) cultivating the cell population under environmental conditions where the gene coding for the cytotoxic polypeptide is expressed, the expression leading to an at least partial killing of the cell population,



(2) a method of containing an extrachromosomal recombinant replicon to a first kind of cell, where said replicon is naturally transferable to a second kind of cell, which method comprises providing on the recombinant extrachromosomal replicon a gene whose expression results in the formation of a cytotoxic polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide, the plasmid F CcdB polypeptide, the plasmid R1 PemK polypeptide, the plasmid RP4 ParE polypeptide, the prophage P1 Doc polypeptide and a functionally equivalent polypeptide for anyone of said polypeptides to an extent whereby the function of the cell is being limited, said first kind of cells having or being modified to have a chromosomal replicon comprising a regulatory nucleotide sequence the gene product of which inhibits the expression of said gene or the cell function-limiting effect of the polypeptide and thereby protects said first kind of cells, said regulatory gene being lacking in said second kind of cell, whereby, if a cell of the second kind receives said extrachromosomal recombinant replicon said gene is expressed and has a function-limiting effect on said second kind of cell, and

15

(3) a method of stochastically limiting in an environment the survival of a cell population, the method comprising transforming the cells thereof with a recombinant replicon containing a regulatably expressible gene which, when expressed in a cell, encodes a cytotoxic polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide, the plasmid F CcdB polypeptide, the plasmid R1 PemK polypeptide, the plasmid RP4 ParE polypeptide, the prophage P1 Doc polypeptide and a functionally equivalent polypeptide for anyone of said polypeptides, the expression of said gene leading to formation of the polypeptide to an extent whereby the function of the cells is being limited, the expression of said gene is stochastically induced as a result of re-combinational excision of an excisable negatively functioning regulatory nucleotide sequence which, while present in the cells, inhibits expression of the gene coding for the polypeptide, said negatively functioning regulatory nucleotide sequence being contained in the recombinant replicon or in an other recombinant replicon present in cells of the population containing the replicon.

25

## DETAILED DISCLOSURE OF THE INVENTION

One objective of the present invention is to provide a novel approach to conditionally controlling the survivability of a recombinant microbial cell population. This approach is based on the use of what is generally referred to as proteic killer systems which have been reviewed i.a. by Jensen et al., 1995. These systems consist of two components, a cytotoxin polypeptide (also referred to herein as a first kind of polypeptide) and a corresponding antitoxin or antidote polypeptide (also referred to herein as a second kind of polypeptide) that by binding to the cytotoxic polypeptide inhibits the toxic effect hereof. A general characteristic of such proteic killer systems is that the antitoxin component in contrast to the toxin component is susceptible to protease degradation, resulting in a decay of the antitoxin polypeptide.

As used herein, the expression "microbial cell" includes any prokaryotic and eukaryotic cells as well as cells of *Archae* species. Thus this expression includes cells of bacterial species, fungal species, animal species including invertebrates, vertebrates, mammals, humans and insects, and plant cells.

Thus, in one aspect of the invention there is provided such a method of conditionally controlling the survivability of a recombinant microbial cell population that comprises as the first step, providing in the cells a gene coding for a cytotoxic first kind of polypeptide, which gene is selected from the gene coding for the *E. coli* K-12 RelE polypeptide and a gene coding for a structurally and functionally equivalent polypeptide and, operably linked to the gene, a regulatable regulatory DNA sequence.

Genes which are structurally and functionally equivalent to *relE* are herein collectively referred to as the *relE* gene family or as *relE* homologues. This group of genes including the *E. coli* plasmid P307 derived *relE* homologue encompasses genes the gene products of which have cytotoxic effects and which, relative to *E. coli* K-12 RelE, have at least 20% such as at least 30% e.g. at least 40% identical and conserved amino acids. The sequences listed in the below Table 1.5 are putative RelE homologues.

Whereas, in accordance with the invention, presently preferred recombinant microbial cells are prokaryotic cells such as Gram-negative and Gram-positive bacterial cells, it has been found that the survivability of other microbial cells such as *Archae*, yeast cells, fungal cells, animal cells including human cells and plant cells and replicons of such organisms can be conditionally controlled using the methods of the present invention.

In the present context, the expression "conditionally controlling" refers to a construction of the microbial cell which permits that the gene coding for the cytotoxic polypeptide can be expressed under certain pre-determined environmental conditions whereas under other such conditions, the gene is not expressed. Hence, the survivability of the microbial cells can be made dependent on certain pre-selected conditions.

In accordance with the invention, the survivability of microbial cells is controlled by the expression in the cells of a cytotoxic polypeptide selected from *E. coli* K-12 RelE polypeptide and a functionally equivalent polypeptide. As used herein, the term "cytotoxic" refers not only to a loss of the ability of microbial cells containing the toxin-encoding gene to remain viable as determined by the capability to propagate in media which, under identical environmental conditions, support unrestricted growth of similar cells not containing the toxin-encoding gene, but also to cells having, as a result of the expression of the polypeptide-encoding gene, a limited cell function, the latter expression denoting that the growth of a cell as manifested i.a. by the synthesis of new cell material and the rate of replication of the cell is decreased.

During the experimentation leading to the invention it was surprisingly found that a range of cytotoxic polypeptides according to the invention have the effect that they inhibit translation of genes. This general effect of RelE polypeptides and functionally equivalent polypeptides appears to represent a hitherto undiscovered mechanism for controlling survivability of cells and thus for containment of such cells or replicons in accordance with the methods of the present invention.

Whereas the recognizable manifestation of such limited cell function may ultimately be cell death, it may also be a reduced cell growth appearing as a reduced rate of replication resulting in a reduced increase of cell numbers within a certain period of time as a

result of an increase of the lag phase and/or of the cell doubling time. Other manifestations may be a relatively increased requirement for one or more nutrient components or a relatively higher susceptibility to detrimental environmental factors such as sub-optimal temperatures or cell damages caused by toxic substances.

5

In the present context, the expression "a functionally equivalent polypeptide" refers to a polypeptide that has substantially the same effect on the survivability of microbial cells as the RelE polypeptide of *E. coli* K-12. As it is shown herein, a variety of Gram-positive and Gram-negative bacteria and *Archae* organisms comprise DNA sequences  
10 showing homology with the RelE polypeptide. To the extent gene products of structural homologues of the *relE* gene product show an effect on microbial cell survivability as it is defined above, they are encompassed by the present invention. It will also be appreciated that the term "functional equivalent" includes variants or derivatives of any of the above first kind of polypeptides, the sequences of which have been modified  
15 by substitution, deletion or addition of one or more amino acids and the gene product of which has retained at least part of the cytotoxic function of the gene product of the non-modified sequence.

In the above method, a regulatable regulatory DNA sequence is operably linked to the  
20 gene coding for the cytotoxic polypeptide. In accordance with the invention, such a regulatory sequence can be one with which the gene coding for the cytotoxic polypeptide is naturally associated or it can be a sequence with which the gene is not naturally associated. In the present context, the term "regulatory DNA sequence" is intended to indicate a DNA sequence which directly or indirectly regulates the expression  
25 of the gene coding for the cytotoxic polypeptide at the level of transcription or at the level of translation or at the level of protein function. The regulatory DNA sequence may thus be one, the function of which results in a suppression or inhibition of the activity of the regulatable promoter.

30 Such regulatory DNA sequences are referred to herein as "negatively functioning regulatory DNA sequences". One interesting example of such a regulatory DNA sequence is a sequence coding for a repressor substance which represses the expression of the gene coding for cytotoxically active polypeptide and which substance may, when a cell containing it is released to a human or an animal body or to the outer envi-

ronment where the substance is no longer being expressed, undergo a decay whereby the repression of expression of the cytotoxin-encoding gene is gradually reduced and eventually, when the decay of the repressor is completed, the repression is removed.

- 5 Another example of such a regulatory DNA sequence is a sequence encoding a polypeptide that acts as an antidote or antitoxin for the cytotoxic polypeptide. Such a sequence include the *relB* gene derived from the *relBE* operon of *E. coli* K-12 which is capable of binding to the RelE polypeptide and thereby inhibiting its effect. As also shown herein, sequences encoding such antitoxins can be found in Gram-negative and  
10 Gram-positive bacteria and in *Archae*. Such homologues of the *relB* sequence are encompassed by the present invention.

- In preferred embodiments of the invention, the regulatory DNA sequence may be present in the cell in one or more recombinant replicons and it may be contained in the  
15 same replicon as that containing the cytotoxin-encoding gene or in a different recombinant replicon.

- One way whereby the expression of the cell function-limiting cytotoxic polypeptide can be regulated is by providing in the cell a gene coding for the polypeptide, which  
20 gene is regulated at the level of transcription. The regulation at the level of transcription may be carried out in various ways including a regulation by means of a promoter, regulated by one or more factors. These factors may either be ones which by their presence ensure expression of the gene coding for polypeptide or may, alternatively, be factors which suppress the expression of the gene so that their absence causes the  
25 polypeptide to be expressed.

- Factors regulating the activity of the promoter as defined above may be selected from a variety of factors. Thus, the expression of the gene encoding the polypeptide may be determined by the environmental conditions, by the physiological state of the cells,  
30 or by a cyclical or stochastic event. In the present context, the term "cyclical event" is understood to mean a cyclically recurrent event causing changes in certain factors known to be potentially useful in affecting the expression of genes such as temperature conditions, changes in light intensity or hormonal changes. The term "physiologi-

cal state of the cells" denotes factors such as cell density or the growth phase of cells.

In accordance with the invention, advantageous promoter regulating factors are readily  
5 regulatable factors including the presence or absence of a certain chemical substance  
in the environment or the physical conditions in the environment such as the prevailing  
temperature or other physical factors (e.g. the intensity of the light in the environ-  
ment). Thus, it is possible to envisage containment systems as presently claimed, in  
which the gene coding for the cytotoxic polypeptide is expressed when a certain  
10 chemical substance present in a first environment such as the fermentation medium in  
which the cell is propagated, is not present in a second environment to which the cell  
is released, or when a factor required for the growth or survival of the cell is no longer  
present, or the factor is one which, when it is depleted or exhausted from an envi-  
ronment of the cell, has the desired effect, viz. that the gene is expressed.

15

The promoter regulating the transcription of the gene coding for the cytotoxic poly-  
peptide can also become activated in a second environment of the cell by a chemical  
substance which is not present in a first environment of the cell, but which is present  
in the second environment in sufficient quantities to activate the promoter. Similarly,  
20 the promoter may be a promoter which is activated by a shift in temperature, such as  
a shift from a higher temperature in a first environment as e.g. a fermentation vessel,  
to a lower temperature prevailing in an outside second environment, or the intensity of  
light, in that the promoter may be one which is activated in the presence of light of  
sufficient intensity, but is inactive in the darkness prevailing in a first environment  
25 such as a fermentation vessel.

Where microbial cells as defined herein are cells that are to be released to the outer  
environment in a controlled manner, e.g. to a restricted area of land or to the intestinal  
tract of a human or an animal, the regulatable promoter may be one which is regulated  
30 chemically, i.e. by the presence or absence of a certain chemical substance in the en-  
vironment of the cells as it has been explained above.

However, the regulatable promoter is advantageously a promoter which is activated  
cyclically, e.g. by changes of the temperature, or by a stochastic event. The term

"stochastic event" as used herein is intended to denote an event which occurs at random at a certain frequency per cell per generation or frequency per unit time which, in accordance with the invention, may result in a limitation of the function of the cells in which the activation of expression of the cytotoxic polypeptide occurs, optionally to an extent which leads to the death of the cells. The stochastic event may be occasioned by periodic inversions of the region carrying the promoter or by recombinational excision of a recombinationally excisable negatively functioning regulatory DNA sequence as defined above.

10 It should be noted that in order to ensure a general applicability of the present invention, the promoter used to initiate transcription of the gene coding for the toxic polypeptide is preferably a promoter which is capable of causing expression of said gene in a wide range of cells.

15 In case of regulatable transcription of the polypeptide, the regulatory DNA sequence may e.g. be a promoter isolated from bacterial operons involved in the biosynthesis of amino acids or from bacterial genes, the transcription of which is activated late in the stationary growth phase or from bacterial genes involved in the synthesis of cell surface structures such as fimbriae. Examples of suitable promoters include the *E. coli trp* promoter which becomes activated in the absence of tryptophan, the bacteriophage  $\lambda$   $P_R$  and  $P_L$  promoters controlled by temperature sensitive regulatory DNA sequences, the *Bacillus subtilis* sporulation gene promoters which are activated during sporulation, and the *E. coli* and *Salmonella* fimbriae gene promoters which are activated stochastically.

25

In case of chemically regulatable promoters, the chemical substance, the presence or absence of which determines the activation of the promoter, is suitably selected from carbon or nitrogen sources, metabolites, amino acids, nucleosides, purine or pyrimidine bases or metal ions. When the chemical substance is one which, when present, suppresses promoter activity, it is preferably a substance which rarely occurs in the natural environment in such concentrations that the promoter would not be activated when the cell is released to the natural environment. One example of such a promoter is the *trp* promoter which is repressed in the presence of tryptophan in the environment of the cell, but which is derepressed in the absence of sufficient amounts of

30

tryptophan in the environment. A containment system according to the invention using the *trp* promoter or another promoter being regulated in the same manner, might therefore comprise an amount of tryptophan in a first environment, such as a fermentation vessel, which is sufficient to repress the promoter in such an environment, the  
5 promoter, however, being derepressed when the cell is released from the first environment to a second environment, e.g. the outer environment which usually contains very low amounts of tryptophan or no tryptophan at all.

It is also possible to select a promoter that is regulated by the absence or presence of  
10 one or more compounds in exudates of plants colonized with a recombinant organism according to invention.

In this context, another useful promoter is an arabinose inducible promoter including that contained in the plasmid pBAD (Guzman et al., 1995). Without arabinose added  
15 to the growth medium, the pBAD promoter is completely turned off. However, in the presence of arabinose, strong transcription is induced. This particular promoter is repressible by the addition of glucose to the growth medium. Thus, by the addition of glucose, transcription from pBAD can be rapidly and efficiently turned off. The glucose repression effect is epistatic to the inducer effect by arabinose. Hence, if cells with a  
20 pBAD-carrying plasmid are grown in a medium containing both arabinose and glucose, the promoter is not induced. However, if cell growth depletes the medium for glucose, then the promoter will be induced. Therefore, such a plasmid is suitable for the conditional turning on and off the expression of gene, in particular toxin-encoding genes as described herein.

25 Accordingly, in one embodiment of the invention the method is used to contain microbial cells wherein the promoter is suppressible by a first kind of chemical compound and inducible by a second kind of chemical compound whereby, when the first kind of compound is depleted from the medium, the promoter is induced by the second kind  
30 of compound.

Another example of a regulatable promoter, the activation of which is determined by a chemical substance is the *lac* promoter which is inducible by e.g. isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG).



As mentioned above, the regulatable promoter may be a promoter, the activity of which is determined by the temperature prevailing in the environment of a cell containing the gene coding for the cell function-limiting cytotoxin and a regulatable promoter regulating the expression of the gene. In such a case, the regulation of the promoter is advantageously obtained by the presence in the cell of a gene coding for a temperature sensitive repressor for the promoter. As one typical example, the  $\lambda$  promoters including those mentioned above may be regulated by a temperature sensitive  $\lambda$  cl repressor.

10

Promoters which are activated stochastically by periodic inversions of the promoter region (in the present context, such promoters are also termed as "invertible promoters" and "inversional switch promoters") and which are useful for the purposes of the present invention include as examples the *hin*, *cin* and *gin* promoters. One particularly useful invertible promoter is the *fimA* promoter which is one *E. coli* fimbriae promoter. The activation (inversional switch) of this promoter is regulated by the gene products of the two genes which for the present purposes is termed the "on" and the "off" genes, the on gene product inducing a switch from off (inactive) to on (active), and the off gene product inducing a switch from on to off. In a wild-type *E. coli* cell where the *fimA* gene and its associated promoter is present in one copy on the chromosome, the inversional switch occurs with a switching frequency of about one cell/1000 cells/generation. It is, however, possible to regulate the frequency of the inversional switch as required by regulating the dosage of expression of the *on* and *off* genes. This is e.g. effected by means of suitable promoters transcribing into the on and off genes. The frequency of transcription initiation by these promoters will then determine the relative dosage levels of the *on* and *off* gene products being formed.

In accordance with the invention, one particular method of stochastically regulating the expression of the gene coding for the toxic polypeptide is the induction of the gene expression as a result of recombinational excision of an excisable negatively functioning regulatory DNA sequence which, while present in the cell, inhibits expression of the gene. In the present context, the term "recombinational excision" refers to the result of a naturally occurring phenomenon of genetic recombination (cross-over) whereby DNA sequences in replicons, in a controlled process, pair, break and rejoin to

30

form recombinant replicons by the sequential action of enzymes acting on the DNA. The frequency of recombinational events in a cell depends i.a. on the degree of homology between paired complementary nucleotide sequences and on the length of the complementary sequences. Thus, it has been shown that about 50 base pairs of homology may be required to obtain recombination in a bacterial cell.

When a negatively regulatory DNA sequence is inserted between directly repeated nucleotide sequences of a sufficient length in a recombinationally proficient cell which, in accordance with the invention contains a gene coding for the toxic polypeptide, recombination between the repeats results in the recombinational excision of the negatively regulatory DNA sequence permitting the gene to be expressed.

Accordingly, the phenomenon of recombinational excision implies that a DNA subsequence, i.e. the negatively regulatory DNA sequence, is excised from a longer DNA sequence through a recombination event. In essence, the longer DNA sequence is cleaved on either side of the subsequence and the fresh ends are joined, leaving out the subsequence. Recombination occurs between sufficient homologous flanking nucleotide subsequences. Thus, with DNA of the general structure W-X-Y-X-Z, X being a repeated sequence and Y being a negatively regulatory DNA sequence, this could recombine to form W-X-Z, with the Y subsequence being excised.

As mentioned above, the frequency of the recombination can be determined by varying the lengths of the repeats and/or the distance between the repeats. Furthermore, the frequency may be varied by using repeat sequences of varying homologies. Thus, nucleotide sequence repeats being 100% homologous and having a size which does not impair recombination will result in a high recombination frequency and hence, in a high frequency of recombinational excision of the negatively regulatory sequence, whereas mismatches within complementary sequences will reduce the recombination frequency depending on the degree of mismatch. As an example, it has been found that 10% divergence between nucleotide sequence repeats may reduce the recombination frequency 40-fold.

Accordingly, the microbial cell comprising the gene coding for the cytotoxic polypeptide may, in accordance with the invention, be a cell containing a regulatory DNA se-

quence which is a recombinationally excisable negatively functioning regulatory DNA sequence being flanked by a first flanking nucleotide sequence and a second flanking nucleotide sequence substantially homologous with the first flanking sequence. As used herein, the term "substantially homologous with" is used to indicate that the degree of homology is sufficient to result in a desired frequency of recombination. In certain embodiments it may, in order to obtain a desirable maximum frequency of recombination, be advantageous to use direct repeats, i.e. sequences being 100% homologous, whereas in other embodiments where a moderate degree of cell function limitation is desirable, it is appropriate to use repeats which are more or less heterologous, but still allowing a desirable lower frequency of recombination to occur. Accordingly, in the present context, the term "sufficiently homologous" is used to indicate a degree of homology between two flanking nucleotide sequence repeats which results in a desired frequency of recombinational events in a cell containing the gene coding for the toxin polypeptide and a negatively regulatory DNA sequence.

15

As it also has been mentioned above, the frequency of recombination depends on the lengths of the flanking sequences. In useful embodiments of the invention, flanking sequences are used which have a length being in the range of 100-5,000 base pairs. In certain preferred embodiments, it is advantageous to use flanking sequences, the length of which is in the range of 200-3,000 base pairs. As the flanking sequences can be used any nucleotide repeats of sufficient lengths and homology as it has been defined above. As one useful example of flanking sequences may be mentioned the chloramphenicol resistance gene having a size of about 900 base pairs and which occurs in the plasmid pBR325. Another example of a useful nucleotide sequence which, when inserted as repeats, results in recombination, is a subsequence of the *rrnB* gene isolated from the plasmid pKK3535 (Brosius et al., 1981, Plasmid, 6:112-118) having a size e.g. in the range of 500 to about 3,000 base pairs, such as 598 base pairs.

In one interesting embodiment of the invention, the excisable negatively regulatory DNA sequence operably linked to the gene encoding the cytotoxic polypeptide is a gene encoding an antisense RNA which forms an RNA-RNA duplex with said the messenger RNA of the polypeptide-encoding gene and thereby, when it is expressed, inhibits translation of said gene coding for the polypeptide.

In another useful embodiment of the present invention, the recombinationally excisable negatively regulatory DNA sequence is a gene encoding a polypeptide repressor of transcription of the polypeptide-encoding gene. Such a repressor may, e.g. be a *lac* repressor including the repressor encoded by the *LacIq* gene. It will be appreciated  
5 that the negatively regulatory DNA sequence can also be a gene coding for RelB anti-toxin or functionally equivalents hereof.

In a further useful embodiment of the invention, the excisable negatively regulatory DNA sequence is a transcription termination sequence, preventing the transcription of  
10 the cytotoxic polypeptide-encoding gene. In one specific embodiment of the invention, such a suitable terminator sequence is the *rpoCt'* transcription terminator isolated from the plasmid pHBA102rpoCt (Squires et al., 1981, Nucleic Acid Res., 9:6827-6839).

Negatively regulatory DNA sequences which, in accordance with the invention, are  
15 suitable, can be isolated from DNA sequences derived from a virus, or a prokaryotic or eucaryotic cell. Thus, sources of the DNA sequence include bacterial chromosomes, bacterial plasmids, prokaryotic viruses, eucaryotic viruses, eucaryotic plasmids, or eucaryotic chromosomes.

20 In preferred embodiments of the invention, the excisable negatively regulatory DNA sequence and the first and second flanking sequences, both as defined above, is provided in the form of a "cassette" which term is used herein to describe a readily insertable DNA sequence comprising at least the above-mentioned sequences and optionally the gene coding for the cytotoxically active polypeptide, and optionally further  
25 nucleotide sequences including as examples a suitable marker such as a gene coding for antibiotic resistance. In the present context, the term "insertable" denotes that the cassette as defined herein is provided with suitable restriction sites at both ends allowing for insertion in a replicon having the same restriction sites. Accordingly, such preferred restriction sites include sites which occur frequently in replicons where inser-  
30 tion is desirable or alternatively, restriction sites which may be easily provided in such replicons.

It will be understood that, in accordance with the invention, a cassette as defined above and which does not comprise the gene coding for toxin polypeptide and opera-

bly linked to the negatively regulatory DNA sequence, may be inserted in a replicon which is different from the replicon containing said gene. Optionally, the cassette as defined above is inserted in a first replicon such as e.g. a transposon and subsequently inserted via the transposon into the chromosome to obtain a cell as defined herein.

5

As it has been explained above, the activation of certain invertible promoters such as the *fimA* promoter or functional homologues hereof is regulated by the gene products of an *on* gene and an *off* gene. It will be understood that this mechanism of promoter regulation provides the possibility of using the *off* gene or a functional homologue

10 hereof as a negatively regulatory DNA sequence which may be inserted in the microbial cell as defined herein, as a recombinationally excisable DNA sequence in the manner explained in details above. Accordingly, in one embodiment, the present invention provides a microbial cell wherein the toxin-encoding gene is stochastically expressed as a result of recombinational inversion of an invertible promoter sequence.

15

In plasmids, inherent mechanisms occur whereby multimer resolution of the plasmid during replication takes place. As exemplified by the broad host range plasmid RP4, this resolution system may comprise (1) a gene coding for a multimer resolving enzyme, a resolvase and (2) a site for the site-specific resolvase-mediated resolution. In  
20 plasmid RP4 the gene coding for the resolvase is *parA* and the site for the resolution is designated *mrs*. If two *mrs* sites are placed in direct orientation, a DNA sequence inserted between those two sites may, if the *parA* gene is present in the same host cell, be deleted at a relatively high frequency whereby a site-specific recombination system is provided. In useful embodiments the *parA* gene may be located in trans.

25

It has been found that such a site-specific recombination system provides a useful mechanism for stochastically regulating the expression of a gene such as the gene coding for the toxic polypeptide as defined herein, since the site-specific recombination may be used to obtain recombinational excision of a negatively regulatory DNA  
30 sequence as defined above.

Accordingly, in one interesting embodiment, the present invention provides a microbial cell as defined herein in which the negatively regulatory DNA sequence is a sequence flanked by a first site for a site-specific resolution recombinase and a second site for

site-specific resolution, the second site being recognizable by the same or a functionally equivalent multimer resolving enzyme as is the first site, whereby the regulatory sequence is recombinationally excisable in the cell. In a specific embodiment, the gene coding for the multimer resolving enzyme is located *in trans* relative to the sites for site-specific resolution. In the present context, one useful example of a suitable gene is the *parA* gene isolated from plasmid RP4.

In accordance with the invention, the method of controlling the survivability of microbial cells can be based on providing in the cells a gene coding for a cytotoxic polypeptide that is structurally and functionally equivalent to the *E. coli* RelE polypeptide (the *relE* gene family). Such a gene can be derived from the chromosome or another replicon of a Gram-negative bacterium including *Enterobacteriaceae* spp. such as *E. coli*, *Hemophilus* spp. such as *H. influenzae*, *Vibrionaceae* spp. such as *V. cholerae*, *Pseudomonadaceae* spp., *Helicobacter* spp. such as *H. pylori* and *Synechosystis* spp, the latter organisms belonging to the group of cyanobacteria. The gene may also be derived from the chromosome and other replicons of Gram-positive bacteria including lactic acid bacteria such as *Streptococcus* spp including *Streptococcus pneumoniae*, *Bacillaceae* spp. such as *B. thuringiensis*, and *Mycobacterium* spp. and from species belonging to *Arhae* such as *Methanococcus jannaschii* and *A. fulgidus*. Such genes include those that are defined herein as belonging to the *relE* gene family. The RelE equivalent polypeptide from *M. jannaschii* was shown to be toxic for *E. coli* when expressed in this organism.

However, genes coding for cytotoxins of other proteic killer systems and which are therefore functional equivalents of the *E. coli* K-12 RelE polypeptide can also be used in accordance with the invention for conditionally controlling the survivability of microbial cells. Such genes include the gene coding for the plasmid F CcdB polypeptide, the gene coding for the plasmid R1 PemK polypeptide, the gene coding for plasmid RP4 ParE polypeptide and the gene coding for the prophage P1 Doc polypeptide, as described by Jensen et al., 1995.

It will be understood that in this context, the term "functional equivalent" includes variants or derivatives of any of the above first kind of polypeptides the sequences of which have been modified by substitution, deletion or addition of one or more amino

acids and the gene product of which has retained at least part of the function of the gene product of the non-modified sequence.

In accordance with the invention, the *relE* family gene or any gene coding for a toxin  
5 of a proteic killer system is provided in the microbial cells at a location where it can be expressed effectively. Thus, in useful embodiments the gene is present on the chromosome of the cells whereas in other embodiments it is preferably located on an extrachromosomal element such as a plasmid or a cosmid. In a specific embodiment, the microbial cells according to the invention do not contain a gene coding for a second  
10 type of polypeptide that is capable of counteracting the cell toxic effect of the RelE polypeptide or the functional equivalent hereof.

However, in other useful embodiments, the microbial cells comprise a gene coding for a second kind of polypeptide that is capable of binding to the *relE* polypeptide or the  
15 functional equivalent hereof, the binding resulting in that the toxic effect of the RelE polypeptide or the functional equivalent is at least partially counteracted. Such a counteracting second kind of polypeptide is, as it is mentioned above, also referred to herein as an antitoxin or an antidote for the cytotoxic polypeptide.

20 Although, in certain uses of the present method, it is preferred that the genes coding for both the toxic polypeptide and the antitoxin herefor is under the control of the same regulatory sequences, it may, in other uses, be advantageous that the gene coding for the second kind of polypeptide is operably linked to a different regulatable regulatory DNA sequence as defined above, permitting that the gene coding for the  
25 second kind of polypeptide is suppressed under conditions where the gene coding for the RelE polypeptide or the functional equivalent is expressed.

It will be appreciated that the genes coding for the toxin polypeptide and the antitoxin polypeptide, respectively can be present on the same replicon such as a plasmid or on  
30 the chromosome, or they can be present on different replicons in the microbial cells.

A useful second kind of polypeptide is the *RelB* polypeptide derived from *E. coli* K-12 which i.a. binds effectively to the *E. coli*-derived RelE polypeptide. However, the regulation of the toxic effect of the first kind of polypeptide can also be based on pro-

viding in the cells a gene coding for a second kind of antitoxically active polypeptide that is functionally equivalent to the *E. coli* RelB polypeptide. Such a gene can be derived from the chromosome or another replicon of a Gram-negative bacterium including *Enterobacteriaceae* spp. such as *E. coli*, *Hemophilus* spp. such as *H. influenzae*,  
5 *Vibrionaceae* spp. such as *V. cholerae*, *Pseudomonadaceae* spp., *Helicobacter* spp. such as *H. pylori*, and *Synechosystis* spp. belonging to the group of cyanobacteria. Additionally, genes coding for structural and functional equivalents of the *E. coli* RelB polypeptide can be isolated from Gram-positive bacteria including lactic acid bacterial species such as *Streptococcus* spp., *Bacillaceae* spp. such as *B. thuringiensis*, and  
10 *Mycobacterium* spp. and from species belonging to *Arhae* such as *M. jannaschii* and *A. fulgidus*. Sequences for the *E. coli* RelB polypeptide and for equivalents isolated from the above organisms are listed in Table 1.6.

Genes coding for functional equivalents of the *E. coli* K-12 RelB polypeptide which in  
15 accordance with the invention can be used for containing microbial cells and replicons include the genes coding for the plasmid F CcdA polypeptide, the plasmid R1 PemI polypeptide, the plasmid RP4 ParD polypeptide and the prophage P1 Phd polypeptide.

It will be understood that in this context the term "functional equivalent" includes  
20 variants or derivatives of any of the above second kind of polypeptides, the sequences of which have been modified by substitution, deletion or addition of one or more amino acids and the gene product of which has retained at least part of the function of the gene product of the non-modified sequence.

25 It is a significant objective of the present invention to provide the means of conditionally controlling the survivability of microbial cells that expresses one or more genes coding for a gene product of interest. In accordance with the invention such an objective is pursued for any type of gene products including enzymes such as proteases, enzymes which are effective in degrading carbohydrates such as starch degrading en-  
30 zymes, lipid degrading enzymes and nucleases.

However, it is of particular interest to provide containment of microbial cells wherein the gene product of interest is selected from an immunologically active gene product,



a gene product that is effective in degradation of an environmental pollutant and a pesticidally active product.

Accordingly, in such specific embodiments the microbial cells are cells which further  
5 comprise a DNA sequence that is selected from a sequence coding for an immunologically active gene product, a sequence coding for a pesticidally active gene product and a sequence coding for a pollutant degrading gene product.

In the present context, the term "immunologically active gene product" is used to describe an epitope (antigenic determinant) from a pathogenic organism which, when it  
10 is administered to the body of a human or an animal, is capable of stimulating the formation of antibodies therein. A microbial cell as defined herein which contains one or more genes encoding such a gene product can be utilized in the preparation of live vaccines. In the immunization against several pathogens it is considered advantageous  
15 to administer live vaccines as compared to killed organisms or antigenic fragments of the pathogen, since the level of immunity conferred by a live vaccine is frequently higher than that conferred by vaccines comprising killed pathogenic organisms or fragments thereof. Most currently used vaccines comprising viable epitope-containing organisms are either based on recombinant non-pathogenic organisms encoding the  
20 epitope or they are based on attenuated pathogenic organisms. The cell advantageously contains a multiplicity of genes each of which codes for a specific immunologically active gene product.

However, up till now the use of live vaccines has been limited since it is difficult to  
25 obtain the right combination of attenuation, viability and adequate immune response. Furthermore, the deliberate release of genetically engineered microorganisms to the body and to the external environment which is a result of the use of viable recombinant organisms as vaccines, is currently not allowed in any country for reasons of public concern as to the possible long-term environmental impact, in particular the risk  
30 of permanent establishment of the GEMs in the environment.

The present invention provides advantageous means of circumventing these problems associated with the use of known GEM-based live vaccines by introducing into a viable epitope-containing cell the regulatably expressible gene coding for a cell toxic

polypeptide as defined above. In particularly interesting embodiments, the invention provides, as a useful basis for a viable vaccine, the microbial cells as defined above whose expression is stochastically induced.

- 5 In useful embodiments of the invention, the cell which contains the DNA sequence coding for an immunologically active gene product further comprises means for transporting the epitope, when expressed, to the outer surface of the cell, i.e. translocating it across the cell membrane. Preferably such a translocation is obtained by inserting the gene coding for the epitope into a nucleotide sequence coding for an outer cell
- 10 surface polypeptide structure such as fimbriae which contains the fimbrillin protein, pili, flagellae or certain other surface proteins including as an example the OM protein found in *Streptococcus* species. By providing the cell with such a hybrid nucleotide sequence being expressible in the cell, the gene product hereof will be a fusion or hybrid protein comprising the epitope and the relevant cell surface structure.

15

- A cell in which a fusion protein is expressed which comprises the epitope fused to a surface structure protein by which the cell can adhere to the mucosal cells of a body to which the cell is administered is considered to be particularly useful in that the epitope will become in close contact with the mucosa and thereby effectively stimulate a
- 20 protective immune response in the form of the excretion of secretory antibodies of the IgA and IgG classes.

- Furthermore, the adhesion of the epitope-carrying cell will ensure that the cell is retained in the human or animal body for a period of time which is sufficient to obtain
- 25 the desired immune response. It is considered that a satisfactory immunization typically may be obtained if the cell is present in sufficient numbers in a particular body environment such as the intestinal tract for a period in the range of 15-30 days, depending on the nature and the activity of the epitope expressed from the cell.

- 30 As it will be understood from the above description of the gene coding for the cell function-limiting toxic polypeptide and the DNA sequence regulating its expression, the present invention may provide useful means of providing live vaccines based on recombinant organisms which are immunologically effective and which can be used

without the risk of undesired spreading of recombinant genes to the microflora of humans and animals or to the outer environment.

In accordance with the invention, a useful cell for the preparation of a live vaccine is one selected from a bacterial species which inherently contains an outer surface structure as mentioned above. Such species include as examples species of *Enterobacteriaceae* such as *Salmonella* and *E. coli* species, *Vibrionaceae* and *Pseudomonadaceae*. It will be understood that strains of such species which are particularly useful in the present invention as the basis of a live vaccine as defined above, are non-pathogenic strains or strains having a low pathogenicity.

The epitope expressed by a cell as defined above may be an epitope derived from any pathogenic organism or agent the obtainment of immunity against which is desirable. Such pathogens include viruses, bacteria and eukaryotic organisms such as fungi, yeast or protozoa.

In commercially important embodiments, the microbial cell comprising the gene coding for a cytotoxic polypeptide contains a nucleotide sequence coding for a pesticidally active gene product. In this context, the term "pesticidally active gene product" is used to denote a product which, when expressed in a cell being released to an environment where there is a need to reduce or eliminate the presence of pests that feed on plants, including insect pests, nematodes and vermins such as rodents or birds, is effective in respect of such pest control.

Such pests are currently controlled by the administration of toxic chemical pesticides to the infested environment, but recently various naturally occurring pesticidally active organisms including viruses, bacteria and fungi have been used as biological pest control products.

Prominent examples of such pesticidally active organisms include biotypes or strains of the species *Bacillus thuringiensis* that produce crystalline proteins being toxic to insects, in particular to caterpillars, and several viruses being pathogenic for insects in the larval stage or in the adult stage. However, the pesticidal effect of such organisms is frequently less satisfactory and there is a strong need in farming, forestry and horti-

culture to provide improved pesticidally active organisms. One approach to solving this problem is to construct genetically engineered organisms having an increased toxic effect or a better survival rate in the environment. In addition to pesticidally active compounds from *B. thuringiensis*, such compounds are produced by other microbial  
5 organisms including *Bacillus sphaericus*, fungal species, algal species and plants. In accordance with the invention, genes coding for such biopesticides can be inserted and expressed in the biologically contained cells of the invention.

To the extent such improved organisms are developed, their use in the environment  
10 will, as a consequence of current public concern of the potential risks involved in deliberate release of such toxic or pathogenic GEMs, only be approved by official environmental agencies if it can be demonstrated that the release does not lead to an undesired propagation or to an extended survival of such organisms in the environment to which they are applied.

15

The present invention clearly provides the means of limiting the survival in the environment of genetically engineered pesticidally active organisms. As it has been explained above, the rate of expression of the cytotoxic polypeptide can be regulated stochastically and thus the survival rate of pesticidally active cells may conveniently be  
20 adapted to any specific need. Also, the cell function-limiting effect of the toxic polypeptide may, in accordance with the present invention, be adjusted by selecting a first kind of polypeptide that has an appropriate cell function-limiting effect.

In another useful embodiment, the invention provides a cell in which the gene coding  
25 for a desired gene product is a sequence coding for a pollutant-degrading gene product. It is known that several xenobiotic compounds polluting the outer environment including soil and water can be degraded by microorganisms having an inherent capability of degrading these compounds. Obviously, the technology of genetic engineering provides means of providing improved organisms having an increased pollutant-de-  
30 grading capacity or having the capacity to degrade a broad range of compounds, in particular hydrocarbons.

However, the public concern as mentioned above are also relevant in this context and accordingly, the present invention provides useful means of providing improved pollu-

tant-degrading microbial cells, the survival of which can be controlled by regulating the expression of the first kind of polypeptide as it is defined above. In particularly preferred embodiments, the cell contains a gene coding for a pollutant-degrading gene product, the expression of which is induced by the presence of a pollutant degradable  
5 by the cell.

In addition to the above desired gene products, the microbial cells according to the invention can express any desired gene product including pharmaceutically active products such as e.g. hormones, interleukines and antibioticly active peptides.

10

As mentioned above, the invention provides in a further aspect a method of confining an extrachromosomal replicon to a microbial cell population. Basically, the method comprises the steps of isolating or constructing a microbial cell containing a gene belonging to the *relE* gene family expressing a first kind of polypeptide that is toxic for  
15 the cell and introducing into the cell the extrachromosomal replicon to be confined, which replicon contains a gene coding for a second kind of polypeptide acting as an antitoxin for said first kind of polypeptide, and cultivating the cells under conditions where the genes coding for the first and the second kind of polypeptides are expressed, whereby a daughter cell that does not receive a copy of the extrachromo-  
20 somal replicon is killed by the first kind of polypeptide being expressed in the absence of expression of the second kind of polypeptide.

In preferred embodiments of such a method the cell population consists of cells that comprise a gene coding for a gene product of interest as defined above.

25

The above method of confining an extrachromosomal replicon is particularly useful when the replicon is a plasmid that naturally occurs in a host cell in a low copy number. Accordingly, the method is useful for confining a plasmid occurring in the microbial cells at a copy number which is in the range of 1-30 including the range of 1-10  
30 such as the range of 1-5.

Microbial cells to which a replicon can be confined in accordance with the invention include Gram-negative bacterial species such as species belonging to *Enterobacteriaceae*, *Hemophilus*, *Vibrionaceae* and *Pseudomonadaceae* and Gram-positive bacterial

species, fungal cells including yeast cells, animal cells including human cells and insect cells, and plant cells.

In a still further aspect, the invention provides a method of post-segregationally stabilizing a plasmid in a microbial host cell population as described above. As it is mentioned above, the method comprises the steps of (i) inserting into the plasmid a gene coding for a first kind of polypeptide as defined herein and a gene coding for a second kind of polypeptide as also defined herein that is capable of being degraded in the host cell at a higher rate than that at which the first kind of polypeptide is degraded, (ii) cultivating the cell population under conditions where the genes coding for the first kind and second kind of polypeptides are expressed, whereby a daughter cell that does not receive at least one copy of the plasmid is killed as a result of the faster degradation of the second kind of polypeptide.

The invention also provides a recombinant microbial cell as defined above, comprising a gene coding for a first kind of polypeptide. Such a cell can be a bacterium of a Gram-negative bacterial species including *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp. and *Pseudomonadaceae* spp or it can be of a Gram-positive bacterial species such as a *Bacillus* species or lactic acid bacterial species, a fungal cell including a yeast cell, an animal cell including a human cell and an insect cell, and a plant cell.

As also mentioned above, the invention pertains in another aspect to a method of limiting the survival of a cell population in a first or a second environment, which method comprises as the first step that the cells are transformed with a gene coding for a cytotoxic polypeptide, which gene is selected from the group consisting of the gene coding for the *E. coli* K-12 RelE polypeptide, the gene coding for the plasmid F CcdB polypeptide, the gene coding for the plasmid R1 PemK polypeptide, the gene coding for plasmid RP4 ParE polypeptide, the gene coding for the prophage P1 Doc polypeptide and a gene coding for a functionally equivalent polypeptide for anyone of said polypeptides.

In a specific embodiment of such a method, the survival of the cell population is limited in a first environment in which the gene is expressed whereby the cell population

is contained in said first environment. In another embodiment, the survival of the cell population is not limited when present in a first environment, which first environment could change to a second environment physically and/or chemically distinct from the first environment, in which first environment the gene whose expression results in the formation of a cytotoxically active polypeptide is not expressed, but the survival of which cell population is limited when transferred to a second environment or when present in a physically and/or chemically changed first environment, where the gene is expressed.

- 10 In a still further embodiment of the above method, the survival of a cell population is being limited by providing in the cells a gene coding for a cytotoxic polypeptide which is operably linked to a DNA sequence encoding an antitoxin repressor substance which can undergo a decay when said cells are released to the outer environment to an extent whereby the repressor substance is converted to a non-functional form, whereby
- 15 as a result of said decay, the function of the cells of the population will be gradually limited.

- In yet another aspect of the invention, there is provided a method of containing an extrachromosomal recombinant replicon to a first kind of cell, where said replicon is
- 20 naturally transferable to a second kind of cell, which method comprises as the first step providing on the recombinant extrachromosomal replicon a gene whose expression results in the formation of a cytotoxic polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide, the plasmid F CcdB polypeptide, the plasmid R1 PemK polypeptide, the plasmid RP4 ParE polypeptide, the prophage P1 Doc
- 25 polypeptide and a functionally equivalent polypeptide for anyone of said polypeptides.

- In one specific embodiment of such a method the gene product which inhibits the expression of the expression of the gene coding for the polypeptide or the cell function-limiting effect of the polypeptide is selected from the *E. coli* relB polypeptide, the
- 30 plasmid F CcdA polypeptide, the plasmid R1 PemI polypeptide, the plasmid RP4 ParD polypeptide, the prophage P1 Phd polypeptide and a functionally equivalent polypeptide of anyone of such polypeptides.

The invention also provides a method as defined above of stochastically limiting in an environment the survival of a cell population. Such a method is particularly useful in the containment of recombinant cells which are to be released to the outer environment or the animal or human body.

5

The invention will now be described in further details in the following examples and the drawings wherein

Fig. 1 illustrates *relB<sub>K-12</sub>::lacZ* and *relE<sub>K-12</sub>::lacZ* translational fusions. Shown are relevant parts of the *lacZ* reporter plasmids pKG4001 (carrying a *relB<sub>K-12</sub>::lacZ* fusion) and pKG4002 (carrying a *relE<sub>K-12</sub>::lacZ* fusion). Numbers to the right in the Figure indicate *lacZ* expression levels in Miller units. The low expression level of *relE::lacZ* in pKG4002 is, in part, due to the presence of an intact *relB* gene located on the plasmid. The *relB* gene product represses the *relBE* promoter c. 130-fold;

15

Fig. 2 illustrates in vitro translation of *relBE<sub>P307</sub>*-carrying plasmids. Lane 1: pBR322; lane 2: pHA402 (pBR322-*relB*<sup>+</sup>); lane 3: pHA403 (pBR322-*relBE*<sup>+</sup>); lane 4: pBR322; lane 5: pHA100 (pBR322-E11 contains the P307 *relBE* genes in their natural context); lane 6: pKG325; lane 7: pHA110 (pBR325-*relB*<sup>+</sup>);

20

Fig. 3 shows the structure of expression plasmid pNDM220. The plasmid is a mini-R1 vector whose copy number is amplifiable at 42°C due to the insertion of the temperature inducible  $\lambda$  P<sub>R</sub> promoter upstream of the replication control region. The plasmid also carries the *cl<sub>857</sub>* temperature-sensitive allele of the *cl* repressor. Genes shown are *copB* (copy number control), *repA* (initiation of replication), *parM* and *parR* (plasmid stability loci), *b/a* ( $\beta$ -lactamase) and *lac<sup>R</sup>*. The plasmid contains the *LacI* regulated pA1/O4/O3 promoter upstream of a multiple cloning site that contains unique *BamHI* and *EcoRI* restriction sites. Thus genes inserted downstream of the promoter are inducible with IPTG;

30

Fig. 4 illustrates cell killing by *relE<sub>K-12</sub>* and anti-killing by *relB<sub>K-12</sub>*. Shown are optical density at 450 nm and viable counts as function of time for strains MC1000/pMG223 (*relE*<sup>+</sup>) (A, B), MC1000/pMG223/pMG2201 (*relB* control plasmid) (C, D) and MC1000/pMG223/pMG2202 (*relB*<sup>+</sup> plasmid) (E, F). At time zero, transcription of *relE*



on plasmid pMG223 was induced by the addition of IPTG (1 mM). Filled symbols indicate that IPTG was added. As seen from (E) and (F), the presence of *relB* on a second plasmid counteracted *relE* mediated cell killing;

- 5 Fig. 5 shows the structure of expression plasmid pBAD33. The plasmid is a medium copy number pACYC-derived vector. The plasmid carries the arabinose inducible pBAD-promoter and the *araC* gene of *E. coli*. Thus upon addition of arabinose to pBAD33 containing cells, genes inserted downstream of pBAD are transcriptionally induced. Genes shown in the Figure are: pACYC-ori: origin of replication; CM(R): gene  
10 encoding chloramphenicol acetyl transferase; bla': truncated (nonfunctional) gene encoding  $\beta$ -lactamase; mRNA1 encodes AraC activator protein; pBAD: arabinose-inducible promoter;

- Fig. 6 A/B illustrates cell killing by  $RelE_{P307}$  and anti-killing by  $RelB_{P307}$ . Shown are optical density at 450 nm (A,C) and viable counts (B, D) as a function of time for strains  
15 MC1000/pHA810/pBR322 (A, B) or MC1000/pHA810/pHA110 (carrying *relB*<sub>P307</sub>). At time zero, transcription of *relE*<sub>P307</sub> on plasmid pHA810 was induced by the addition of arabinose (0.02%). Filled symbols indicate that arabinose was added. As seen from (C) and (D), the presence of *relB*<sub>P307</sub> on a second plasmid counteracted *relE*<sub>P307</sub> mediated cell killing;  
20

Fig.7 shows maps of pHA705 and pHA715;

- Fig. 8 illustrates OD<sub>450</sub> of MC1000/pHA-Sp2, MC1000/pHA705 and MC1000/pHA715 (+/- IPTG);  
25

Fig. 9 shows viable counts of MC1000/pHA-Sp2, MC1000/pHA705 and MC1000/pHA715 (+/- IPTG);

- Fig. 10 is the DNA sequence of the *relBE*<sub>Sp2</sub> locus of *S. pneumoniae*;  
30

Fig. 11 is a map of pHA-Sp2;

Fig. 12 is a map of pHAG33;

Fig. 13 is a map of pHAG33-2;

Fig. 14 is a map of pHAG33-3;

5 Fig. 15 is a map of pHAG33-4;

Fig. 16 illustrates OD<sub>450</sub> of KT2440/pHAG33-2, KT2440/pHAG33-3 and  
KT2440/pHAG33-4 (+/- IPTG);

10 Fig. 17 shows viable counts of KT2440/pHAG33-2, KT2440/pHAG33-3 and  
KT2440/pHAG33-4 (+/- IPTG);

Fig. 18 is a map of pHA810;

15 Fig. 19 illustrates Glucose run-out, OD<sub>450</sub> of MC1000/pHA810;

Fig. 20 illustrates Glucose run-out, viable counts of MC1000/pHA810; and

Fig. 21 illustrates that RelE<sub>K12</sub>, RelE<sub>P307</sub> and RelE<sub>Mj</sub> inhibit translation *in vitro*

20

## EXAMPLES

### Materials and methods

25

#### **(i) Bacterial strains**

The *E. coli* K-12 strain MC1000 (Casadaban and Cohen, 1980) which contains a  
chromosomal copy of the *relBE* genes was used as the standard cloning strain and  
30 when a chromosomal copy of the *relB* operon was required. The *E. coli* K-12 strain  
JS115 (*leu*, *thy*, *thi*, *supE*,  $\Delta$ *relB*), which contains a deletion covering the entire *relB*  
operon was provided by Olle Karlström. The latter strain was used for the regulatory  
studies of *relBE*.

**(ii) Plasmids used**

Plasmid pOU253 is a mini-R1 based translational fusion vector carrying the *lacZ* gene of pNM482 (Minton, 1984). The fusion vector is segregationally stable due to the presence of the *parA* system of plasmid R1 (Dam and Gerdes, 1994).

Plasmid pNDM220 is a low copy-number mini-R1 expression vector carrying a multiple cloning site (mcs) placed between the LacI regulated pA1/O4/O3 promoter (Lanzer and Bujard, 1988) and two transcriptional terminators.

pNDM220 was deposited on 30 April 1998 under the Budapest Treaty with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH under the accession No. DSM 12157.

Plasmid pBD2430 (+ 388 - + 1899) is a pUC18 derivative carrying the complete *relBE* operon and gene *IV* located downstream of *relF* (Olle Karlström, unpublished). The relevant *E. coli* DNA present in pBD2430 is shown in Table 1.1 below.

pBD2430 was deposited on 30 April 1998 under the Budapest Treaty with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH under the accession No. DSM 12161.

**(iii) Plasmids constructed**

pKG4001: pBD2430 was digested with *EcoRI* and *XhoI* and the fragment carrying the *relB* promoter (Table 1.1) was inserted into pOU253 producing an in-frame translational fusion between *relB*<sub>K-12</sub> and *lacZ*. Thus, pKG4001 carries a *relB*<sub>K-12</sub>::*lacZ* translational fusion.

pOU253 was deposited on 30 April 1998 under the Budapest Treaty with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH under the accession No. DSM 12158.

pKG4002: pBD2430 was digested with *EcoRI* and *Bst*1107I and the resulting fragment was inserted into pOU253 producing an in frame translational fusion between *relE<sub>K-12</sub>* and *lacZ*. Thus pKG4002 carries an intact *relB<sub>K-12</sub>* gene and a *relE<sub>K-12</sub>::lacZ* translational fusion.

5

pMG223: *relE<sub>K-12</sub>* was amplified by PCR on pBD2430 with primers *relE1B* (5'-CCCCGGATCCATAAGGAGTTTTATAAATGGCGTATTTTCTGGATTTTGACG, SEQ ID NO:1) containing the *parA* Shine & Dalgarno (Gerdes and Molin, 1986) and *relE2* (5'-CCCCCCTCGAGGTCGACTCAGAGAATGCGTTTGACCGC-3', SEQ ID NO:2). The resulting *relE<sub>K-12</sub>* carrying fragment was inserted into pNDM220 using the *Bam*HI and *Sal*I restriction sites. Plasmid pMG223 expresses RelE<sub>K-12</sub> upon addition of IPTG.

pMG2201: this plasmid contains the *EcoRI-Eco*47III fragment from pBD2430 inserted between the *EcoRI* and *Scal* sites of pBR322. Plasmid pMG2201 carries the *relB<sub>K-12</sub>* promoter and the 5' part of the *relB<sub>K-12</sub>* gene.

15

pMG2202: pBD2430 was digested with *EcoRI* and *Bst*1107I and the *relB<sub>K-12</sub>*-carrying fragment was inserted into pBR322 *EcoRI-Scal*. The resulting plasmid carries the *relB* promoter and *relB<sub>K-12</sub>*.

20

pHA100: Plasmid pNZ945 is a pBS(+) derivative that carries a 4.3 kb *EcoRI* fragment from plasmid P307. This fragment encodes the RepFIB replicon and the *relBE* genes of P307 (Saul et al., 1989). The 4.3 kb *EcoRI* fragment (designated E11) of pNZ945 was purified and restricted with *Pst*I. The resulting 2.2 kb *EcoRI-Pst*I fragment was inserted into pBR322 restricted also with *EcoRI* and *Pst*I. The pBR322-derived plasmid carrying the 2.2 kb *EcoRI-Pst*I fragment was designated pHA100. Plasmid pHA100 codes for the entire *relBE* system from P307.

25

pNZ945 was deposited on 30 April 1998 under the Budapest Treaty with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH under the accession No. DSM 12160.

30

pHA110: The 2.2 kb *EcoRI-Pst*I fragment of pHA100 was purified and digested with *Apo*I (*EcoRI* isoschizomer). The resulting *EcoRI-Apo*I DNA fragment (+1 to +1122)

was inserted into the *EcoRI* site of pKG325 which was constructed as follows: Plasmid pBR325 was restricted with *PstI*, which has a unique recognition site in the plasmid. The resulting vector DNA fragment was made blunt ended with T4 DNA polymerase according to the manufacturer's instructions, and religated. Transformants  
5 that were resistant to chloramphenicol and tetracycline, but sensitive to ampicillin were selected. Thus, pKG325 is a TcR, CmlR and ApS derivative of pBR325.

pKG325 was deposited on 30 April 1998 under the Budapest Treaty with the DSMZ-  
Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH under the accession  
10 No. DSM 12159.

Plasmid pHA110 contains the *relB* promoter (*preIB<sub>P307</sub>*) and gene *relB<sub>P307</sub>*.

pHA205: Plasmid pHA205 is a derivative of the low copy-number mini-R1 expression  
15 vector pNDM220 that contains the *relB* gene from P307. The PCR fragment generated from pNZ945 using primers RelB-P307/1: 5'-

CCCCCGGATCCCAGTCTTGAAAGGTGGC-3' (SEQ ID NO: 3) and RelB-P307/2: 5'-

CCCCGAATTCTCATAGGTATTTATCCAG-3' (SEQ ID NO:4) was restricted with

*Bam*HI and *Eco*RI and inserted downstream of the pA1/04/03 promoter of pNDM220.

20 pHA210: Gene *relE<sub>P307</sub>* was PCR-amplified from pNZ945 with the primers: *relE*-p307/3

(5'-CCCCGGATCCAGATCTGGATAAATACC, SEQ ID NO:5) and *relE*-P307/2 (5'-

CCCCGAATTCGTAACCTTTCTGTGTTTATTGC, SEQ ID NO:6). The resulting PCR DNA

fragment was restricted with *Bam*HI and *Eco*RI and inserted into pNDM220 also re-

stricted with *Bam*HI and *Eco*RI. Plasmid pHA210 (+ 1089 to + 1417) is thus a mini-R1

25 derivative carrying a pA1/04/03::*relE<sub>P307</sub>* gene fusion which renders *relE<sub>P307</sub>* inducible with IPTG.

pHA215: Genes *relBE<sub>P307</sub>* were PCR-amplified from pNZ945 with the primers RelB-

P307/1 (5'-CCCCCGGATCCAGTCTTGAAAGGTGGC, SEQ ID NO:3) and *relE*-P307/2

30 (5'-CCCCGAATTCGTAACCTTTCTGTGTTTATTGC, SEQ ID NO:6). The resulting PCR-

generated DNA fragment was restricted with *Bam*HI and *Eco*RI and inserted into

pNDM220 also restricted with *Bam*HI and *Eco*RI. Plasmid pHA215 (+ 840 to + 1417)

is thus a mini-R1 derivative carrying a pA1/04/03::*relBE<sub>P307</sub>* gene fusion rendering the

*relBE<sub>P307</sub>* genes inducible with IPTG.

pHA402: A *Pst*I-*Aat*II fragment from plasmid pHA205, which carries *lac*<sup>R</sup> and the pA1/O4/O3::*relB*<sub>P307</sub> gene fusion was inserted into pBR322 also restricted with *Pst*I and *Aat*II. Thus, the high copy-number plasmid pHA402 contains a *relB*<sub>P307</sub> gene which is inducible with IPTG.

5

pHA403: A *Pst*I-*Aat*II fragment from plasmid pHA215, which carries *lac*<sup>R</sup> and the pA1/O4/O3::*relBE*<sub>P307</sub> gene fusion was inserted into pBR322 also restricted with *Pst*I and *Aat*II. Thus, the high copy-number plasmid pHA403 contains the *relBE*<sub>P307</sub> genes which can be conditionally induced by the addition of IPTG.

10

pHA810: A DNA fragment encoding *relE*<sub>P307</sub> was generated by PCR using primers *relE*-P307/4 (5'-CCCCGAGCTCAGATCTGGATAAATACC, SEQ ID NO:7) and *relE*-P307/5 (5'-CCCCGCATGCGTAACTTTCTGTGTTTATTGC, SEQ ID NO:8). The fragment was digested with *Sac*I + *Sph*I and inserted into the expression plasmid pBAD33 also digested with *Sac*I + *Sph*I. The resulting plasmid, pHA810 (+1089 - +1417), contains the pBAD::*relE*<sub>P307</sub> gene fusion that renders *relE*<sub>P307</sub> inducible with arabinose.

15

An overview of the bacterial strains and plasmids used herein is shown in Table 0.1 below.

20

25

30

Table 0.1. Bacterial strains and plasmids

Strains	genotypes	Reference/Source		
5 MC1000	<i>Δlac leu ara</i>	Casadaban & Cohen, 1980		
JS115	<i>ΔrelB leu thy thi supE</i>	J. P. Bouche, unpublished		
Plasmids	Replicon	Resistance <sup>a)</sup>	<i>relBE</i> co-ordinates <sup>b)</sup>	Reference/Source
10 pOU253	mini-R1	Ap <sup>R</sup>	none	lab. collection
pBAD33	pACYC	Cml <sup>R</sup>	none	Guzman et al., 1995
pNDM220	mini-R1	Ap <sup>R</sup>	none	Gotfredsen & Gerdes, 1998
pBR322	ColE1	Ap <sup>R</sup> , Tc <sup>R</sup>	none	Bolivar et al., 1978
pKG325	pBR325	Tc <sup>R</sup>	none	lab. collection
15 pBD2430	pUC	Ap <sup>R</sup>	+388 - +1899	Olle Karlström collection
pNZ945	pUC	Ap <sup>R</sup>	+1 - +4298	Saul et al., 1989
pKG4001	mini-R1	Ap <sup>R</sup>	+388 - +596	Gotfredsen & Gerdes, 1998
KG4002	mini-R1	Ap <sup>R</sup>	+388 - +921	Gotfredsen & Gerdes, 1998
pHA100	pBR322	Tc <sup>R</sup>	+1 - +2198	Grønlund & Gerdes, 1998
20 pHA110	pBR325	Tc <sup>R</sup>	+1 - +1122	Grønlund & Gerdes, 1998
pHA205	mini-R1	Ap <sup>R</sup>	+840 - +1111	Grønlund & Gerdes, 1998
pHA210	mini-R1	Ap <sup>R</sup>	+1089 - +1417	Grønlund & Gerdes, 1998
pHA215	mini-R1	Ap <sup>R</sup>	+840 - +1417	Grønlund & Gerdes, 1998
pHA402	pBR322	Tc <sup>R</sup>	+840 - +1111	Grønlund & Gerdes, 1998
25 pHA403	pBR322	Tc <sup>R</sup>	+840 - +1417	Grønlund & Gerdes, 1998
pHA810	pACYC	Cml <sup>R</sup>	+1089 - +1417	Grønlund & Gerdes, 1998
pMG223	mini-R1	Ap <sup>R</sup>	+733 - +1020	Gotfredsen & Gerdes, 1998
pMG2201	pBR322	Tc <sup>R</sup>	+388 - +597	Gotfredsen & Gerdes, 1998
pMG2202	pBR322	Tc <sup>R</sup>	+388 - +921	Gotfredsen & Gerdes, 1998

a) Tc<sup>R</sup>, tetracycline resistance; Ap<sup>R</sup>, ampicillin resistance; Cml<sup>R</sup>, chloramphenicol resistance.

b) Co-ordinates refer to Table 1.1 (*relBE*<sub>K-12</sub>, pMG-plasmids) or Table 1.2 (*relBE*<sub>P307</sub>, pHA-plasmids)

#### (iv) Growth media and antibiotics

The growth medium was LB medium (Bertani, 1951) or A + B minimal medium (Clark and Maaløe, 1967) supplemented with 0.2% glucose and 1% casamino acids. For growth on solid media, LA-plates were used. LA is LB containing 15 g agar per litre. All media were supplemented with 50 μg/ml thymine for growth of the strain

JS11507-05-99 $\Delta$ re/BEF<sub>K-12</sub>. Antibiotics were added at the following concentrations: ampicillin, 30  $\mu$ g/ml, and tetracycline, 10  $\mu$ g/ml. When indicator plates were used X-gal (5-Bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside) was added to a final concentration of 40  $\mu$ g/ml.

5

**(v) Conditions of cell growth.**

Cells were diluted in LB + antibiotics from an overnight culture to an OD<sub>450</sub> of 0.005. The cultures were then grown at 37°C until an OD<sub>450</sub> of 0.4 and then diluted to an  
10 OD<sub>450</sub> of 0.01 in 37°C LB containing 1 mM IPTG and antibiotics. Samples for OD<sub>450</sub> measurements and viable counts were taken at the time points indicated. Viable counts were made by plating dilutions of the cultures onto LA plates containing the proper antibiotics.

15 **(vi) Coupled *in vitro* transcription and translation.**

The reactions were performed using the *E. coli* S30 Extract System For Circular DNA as described by the supplier (Promega Corp.). 4  $\mu$ g of DNA was used in all reactions. The reactions were run on a 16% Tricine-SDS-PAGE gel essentially as described by  
20 Schägger and von Jagow (1987).

**(vii)  $\beta$ -galactosidase assays.**

$\beta$ -galactosidase assays were performed essentially as described by Miller (1972).  
25

**(viii) Homology search.**

BLAST searches were performed at the GENESTREAM BLAST network server CRBM Montpellier, France. Standard conditions were used except that the blosum 80 matrix  
30 was used.



## EXAMPLE 1

The occurrence of *relBE* operons in bacteria and *Archae*5 1.1. Nucleotide sequence of the *relBE* operon of *E. coli* K-12

The DNA sequence of the *relBE* operon from *E. coli* K-12 is shown in Table 1.1. In this Table the transcriptional start site of the *relBE* mRNA is indicated with two asterisks (heterogeneity). IR indicates inverted repeats in the promoter and terminator regions.

10 Start codons and stop codons are shown in bold. The transcriptional termination point (*ttp*) of the *relBE* mRNA is also indicated with a vertical arrow. The DNA sequence is from Bech et al., 1985.

By visual inspection of the *relB*<sub>K-12</sub> and *relE*<sub>K-12</sub> genes there was found striking similarity

15 with the so-called "proteic plasmid stabilization systems" as described by Jensen and Gerdes (1995). First, *relE*<sub>K-12</sub> codes for a very basic protein (RelE<sub>K-12</sub>; pI=9.7) of 95 amino acids (aa), and *relB*<sub>K-12</sub> codes for a very acidic protein (RelB<sub>K-12</sub>; pI=4.8) of 79 aa.

20 The sequences of proteins RelB<sub>K-12</sub> and RelE<sub>K-12</sub> are shown in Tables 1.5 and 1.6, respectively. These Tables show multiple sequence alignments of the RelB and RelE gene families. Conserved amino acids at a given position are shown with shading as follows: two amino acids are considered conserved if they both belong to one of the following groups: group 1: D and N; group 2: E and Q; group 3: S and T; group 4: K

25 and R; group 5: F, Y and W; group 6: L, I, V and M. Light grey shading indicates 60-80% conservation, dark grey indicates 80-99% conservation and black indicates 100% conservation. Note in Table 1.6 the fully conserved glycine at position 69 (G in consensus line) and the fully conserved arginine at position 79 (R in consensus line).

The entrez database accession numbers of the protein sequences are given in Tables

30 1.3 and 1.4.

The *relB*<sub>K-12</sub> and *relE*<sub>K-12</sub> genes are co-transcribed with a third gene, *relF* (also denoted *orf-3* or *hokC*), which is homologous to the *hok* gene from plasmid R1 (Gerdes et al., 1986). The start site (i.e. the 5'-end) of the *relBE* mRNA was determined to be 31 nu-

cleotides upstream of the *relB*<sub>K-12</sub> AUG start-codon (Bech et al., 1985) and was confirmed (M. Gotfredsen and K. Gerdes, 1998). Inverted arrows in the *relBE* promoter region (Table 1.1) indicate putative binding sites for regulators of transcription (i.e. the RelB<sub>K-12</sub> and RelE<sub>K-12</sub> proteins themselves).

5

The properties described above suggested that RelE could be a cytotoxin and that RelB could be an antitoxin which counteracts the toxicity elicited by RelE.

Table 1.1. DNA sequence of the *relBE* operon from *E. coli* K-12 (SEQ ID NO:9)

10

```

      1  CTTAATTTCA GGCCCATCG GATCACACAT GGAGAGTTTT TATGAATAAC
    51  CCCGTCTGTC TTGATGACTG GTTGATTGGC TTAAAAGCT TGTGACAGG
   101  GGTAACGTT CGGCAATAAT TTTCTGCCGC ATGCGGGTGT TGCATAAAC
   151  GTGTTACGTT CCTTATCGA CAGGTCAGGT CACCGCTCAC CCGCCGACGA
  201  GAAAGCAACA CTGACATGCT AAAGCAAAAA ATAGATGAAT AAGTTGAGTT

  251  GTGCATATGT AGCCTGACCG TCACAAAGTA TATGGTGTCT GTACCAGTAA
  301  GATGATGGCC GGACTCTTTA AAAACGAGCT GACCTGCACA ATACAGGATG
  401  GACTTAGCAA TGGCTGCTCC TGGCACAAAG CGGACAGTGA TCACCGTTCT
  501  TACGACTACT TTCTGACTTC CTTCTGACT TGCCCTAAGC ATGTTGTAGT

  551  GCGATACTTG TAATGACATT TGTAATTACA AGAGGTGTAA GACATGGGGTA
      **→relBEF mRNA          relB start
      ---->IR<-----

  601  GCATTAACCT GCGTATTGAC GATGAACTTA AAGCGCGTTC TTACGCCGCG
  701  CTTGAAAAAA TGGGTGTAAC TCCTTCTGAA GCGCTTCGTC TCATGCTCGA
  801  GTATATCGCT GACAATGAAC GCTTGCCGTT CAAACAGACA CTCCTGAGTG
  901  ATGAAGATGC TGAACCTGTG GAGATAGTGA AAGAACGGCT TCGTAATCCT

  1001  AAGCCAGTAC GTGTGACGCT GGATGAACTC TGATGGCGTA TTTTCTGGAT
      End relB          Start relE
  1101  TTTGACGAGC GGGCACTAAA GGAATGGCGA AAGCTGGGCT CGACGGTACG
  1201  TGAACAGTTG AAAAAGAAGC TGGTTGAAGT ACTTGAGTCA CCCC GGATTG
  1301  AAGCAAACAA GCTCCGTGGT ATGCCTGATT GTTACAAGAT TAAGCTCCGG
  1401  TCTTCAGGCT ATCGCCTTGT ATACCAGGTT ATAGACGAGA AAGTTGTCGT
  1501  TTTCTGATT TCTGTTGGGA AAAGAGAACG CTCGGAAGTA TATAGCGAGG
      End relE
  1601  CGGTCAAACG CATTCTCTGA ACCAAAGCAT GACATCTCTG TTTCGCACCG

```

Start *hokC* (relF)  
1051 AAGGTGACAC TTCTGCTTTG CGTTGACAGG AGAAGCAGGC TATGAAGCAG  
1101 CAAAAGGCGA TGTTAATCGC CCTGATCGTC ATCTGTTTAA CCGTCATAGT  
5 1151 GACGGCACTG GTAACGAGGA AAGACCTCTG CGAGGTACGA ATCCGAACCG  
End *hokC*  
1201 ACCAGACGGA GGTCGCTGTC TTCACAGCTT ACGAACCTGA GGAGTAAGAG  
10 1251 ACCCGGCGGG GGAGAAATCC CTCGCCACCT CTGATGTGGC AGGCATCCTC  
1301 AACGCACCCG CACTTAACCC GCTTCGGCGG GTTTTGTTT TTATTTTCAA  
15 ----- IR ----- ttp  
1351 CGCGTTTGAA GTTCTGGACG GTGCCGGAAT AGAATCAAAA ATACTTAAGT  
(data base accession number X02405)

Table 1.3. *relE* homologues from Gram-positive and Gram-negative bacteria and *Archae*

	Bacterial species	entrez accession	gene <sup>a)</sup>	Number of aa	MW (kD)	pI
<b>Gram-negative bacteria:</b>						
5	<i>E. coli</i> K-12	132284	<i>relE</i> <sub>K-12</sub>	95	11.2	9.7
10	<i>E. coli</i> K-12	984581	<i>relE</i> <sub>SOS</sub> <sup>b)</sup>	92	10.8	9.5
	<i>E. coli</i> plasmid P307	516611	<i>relE</i> <sub>P307</sub>	95	11.2	9.9
15	<i>H. influenzae</i>	1175293	<i>relE</i> <sub>Hi</sub>	102	11.9	6.7
	<i>V. cholera</i>	396846	<i>relE</i> <sub>Vc</sub>	96	11.2	9.9
	<i>H. pylori</i>	2314031	<i>relE</i> <sub>Hp</sub>	88	10.4	7.9
20	<i>Synechosystis</i>	1653777	<i>relE</i> <sub>Sy</sub>	120	13.7	7.9
<b>Gram-positive bacteria:</b>						
25	<i>B. thuringiensis</i>	520407	<i>relE</i> <sub>Bt</sub>	74	8.6	9.7
	<i>M. tuberculosis</i> #1	2612811	<i>relE</i> <sub>Mt1</sub>	87	10.2	11.0
	<i>M. tuberculosis</i> #2	2695832	<i>relE</i> <sub>Mt2</sub>	97	11.1	9.5
30	<b>Archae:</b>					
	<i>M. jannaschii</i> #1	1498833	<i>relE</i> <sub>Mj1</sub>	90	11.0	10.2
	<i>M. jannaschii</i> #2	1499953	<i>relE</i> <sub>Mj2</sub> (*)	88	10.6	10.0
35	<i>M. jannaschii</i> #3	1591583	<i>relE</i> <sub>Mj3</sub> (*)	91	11.1	10.1
	<i>A. fulgidus</i> #1	2648176	<i>relE</i> <sub>Af1</sub>	87	10.6	10.3
	<i>A. fulgidus</i> #2	2649499	<i>relE</i> <sub>Af2</sub>	92	11.0	9.9
40	<i>A. fulgidus</i> #3	2649496	<i>relE</i> <sub>Af3</sub>	85	10.0	10.0
	<i>A. fulgidus</i> #4	2649514	<i>relE</i> <sub>Af4</sub>	86	10.2	9.9

45 a) *relE* homologues marked with (\*) are not located adjacent to a *relB* partner

b) The *relBE*<sub>SOS</sub> system of *E. coli* K-12 contains a LexA binding-site in the promoter region (Lewis et al., 1994)

Table 1.4. *relB* homologues from Gram-positive and Gram-negative bacteria and *Archae*

5	Bacterial species	entrez accession	gene <sup>a)</sup>	Number of aa	MW (kD)	pI
<b>Gram-negative bacteria:</b>						
	<i>E. coli</i> K-12	132283	<i>relB<sub>K-12</sub></i>	79	9.1	4.8
10	<i>E. coli</i> K-12	984582	<i>relB<sub>SOS</sub></i> <sup>b)</sup>	86	9.4	5.2
	<i>E. coli</i> K-12	984588	<i>relB<sub>K-12,2</sub></i> (*)	97	11.2	5.5
15	<i>E. coli</i> plasmid P307	516610	<i>relB<sub>P307</sub></i>	83	9.2	4.4
	<i>S. typhimurium</i>	731639	<i>relB<sub>St</sub></i> (*)	68	7.6	5.3
	<i>H. influenzae</i>	1573712	<i>relB<sub>Hi</sub></i>	98	11.0	4.7
20	<i>V. cholera</i>	396847	<i>relB<sub>Vc</sub></i>	82	8.9	4.4
	<i>H. pylori</i>	2314037	<i>relB<sub>Hp</sub></i>	95	11.4	9.8
25	<i>Synechosystis</i>	1653776	<i>relB<sub>Sy</sub></i>	86	9.9	4.7
<b>Gram-positive bacteria:</b>						
	<i>B. thuringiensis</i>	520406	<i>relB<sub>Bt</sub></i>	85	10.1	4.5
30	<i>M. tuberculosis</i> #1	2612810	<i>relB<sub>Mt1</sub></i>	93	10.2	4.6
	<i>M. tuberculosis</i> #2	2695833	<i>relB<sub>Mt2</sub></i>	89	9.8	5.1
<b>Archae:</b>						
35	<i>M. jannaschii</i> #1	1498832	<i>relB<sub>Mj1</sub></i>	82	9.6	4.5
	<i>A. fulgidus</i> #1	2648190	<i>relB<sub>AF1</sub></i>	65	7.8	4.8
40	<i>A. fulgidus</i> #2	2649516	<i>relB<sub>AF2</sub></i>	62	7.4	4.3
	<i>A. fulgidus</i> #3	2649510	<i>relB<sub>AF3</sub></i>	72	8.5	4.5
45	<i>A. fulgidus</i> #4	269513	<i>relB<sub>AF4</sub></i>	57	6.7	4.1

a) *relB* homologues marked with (\*) are not located adjacent to a *relE* partner.

b) The *relBESOS* system of *E. coli* K-12 contains a LexA binding-site in the promoter region (Lewis et al., 1994).

1  
55  
58  
55  
56  
52  
54  
54  
54  
54  
58  
77  
72  
59  
71  
58  
55  
32

```

100      *      120      *
:  ---EVA---GDTA---FFRV---HRS---LVDFYP---88
:  K-YE-N-G-KLAFYRV---HR-Q---LYKRF---91
:  --FL-DKPTKTVH-LA---ER---GVAD---87
:  V-LWD---DREI---R---SR-E-G-AVKNP---90
:  E-RNHE---EKVIY-QA---GN---GDYK---74
:  Q-IDEQ---L-LIA-VV---GK-ERSEVAVLASER-MR---95
:  Q-IDEK---V-VVE-ES---GK-ERSEVSEAVKRIL---95
:  Q-ENBI---I-EAV-ES---GK-ERSEVSEAVKALQR-LDD---96
:  R- ---DDEHTTVV-L-ER---DH---ADLYR---87
:  A- ---EDGHRVVE-IL-AR- ---SASIRMNPCPR---97
:  QYIQEFDEIKFSR-LNIHSQTAK---102
:  K-LTSL- ---RFR- TGTH-ALG---92
:  --VVD---DEILR---LGSHSE-LI---88
:  K-N- ---VEDETIF-YSE-HFE-KVA---92
:  D-F- ---LIGNRVV-F-KF-AASE-G-VFKTEKFF---86
:  K-DEETVKF-TFKH---HN-HAYK---85
:  L-SESENS- ---L-LTVSKAEQED-AASDINSILGEYSIED---120

```



### 1.2. Nucleotide sequence of the *relBE* operon of plasmid P307

By database searching it was found that the *E. coli* plasmid P307 codes for a gene system which exhibits both structural and sequence similarity with the *E. coli relBE* genes described above.

The DNA sequence of the *relBE*<sub>P307</sub> genes is shown in Table 1.2. The transcriptional start site of the *relBE* mRNA is indicated with an asterisk, and the -10 and -35 sequence elements of the *relBE* promoter are underlined. The Shine & Dalgarno sequence of the *relB* and *relE* genes are doubly underlined. The DNA sequence is from Saul et al., 1989.

Again, *relE*<sub>P307</sub> codes for a very basic protein of 95 aa (pI = 9.9), and *relB*<sub>P307</sub> codes for a very acidic protein of 83 aa (pI = 4.4), see Tables 1.3 and 1.4. The protein sequences of RelE<sub>P307</sub> and RelB<sub>P307</sub> are also shown in Tables 1.5 and 1.6, respectively. The start site (i.e. the 5'-end) of the *relBE*<sub>P307</sub> mRNA was determined to be located 27 nucleotides upstream of the *relB*<sub>P307</sub> AUG start codon. Inverted arrows in the *relBE*<sub>P307</sub> promoter region (Table 1.2) indicate putative binding sites for regulators of transcription (i.e. the RelB<sub>P307</sub> and RelE<sub>P307</sub> proteins).



Table 1.2. DNA sequence of the *relBE* operon from the *E. coli* plasmid P307 (SEQ ID NO:44)

```

5      301  GAGTATCATA TTAGGATACG GGTGGGTGAC GCCCACCTCT GGCATAGAAC
      351  GGACATTCAT TGATGCCATG CCAGAATGGA CGTTCAGGTT ATTCCGTCCA
      401  GTTCTGCTGG CAACGCGAGA TCTCCCCTGG TATAGTGATG CCACAGCAAA
10     451  GCGCTCAAAC AGGGATAATA TGATGGAAAT CAAGGCTCAA CAGTTTTGTC
      501  ACATCAACGG GGCGGCAAGT CCTTACTGAC AACGGACAAC AAGGTATGGG
      551  CGGCGTGGCG GGTATCGGTT CCACGACTGA AAAGCATCAG GGGCGCGTGG
15     601  CGGAAGCGAT TTTTGCGAAC TGC GCGGAAC TGGATAACGA CCAGCTTAAC
      651  GAGATCATCG AGTGGGTTCG GCTCTATCAG CGCTGAATGC CACTATCAGG
20     701  CTGCGCAAGC GGCCTTTTTT ACGCCCCTTG TTTAATTCCC GCACTACCTG
      751  GACGTTCAGG TGATTCTGTC CATCTGTACA AAAACAATA AAAGACTTGT
                                     -35
25     801  TAACAGGTCA TGTAAGGAGT ATCTTTGAGA CTGGTTAAAC AGTCTTGAAA
                                     -10      *→ relBEP307 mRNA
      851  GGTGGCCTAT GCCTAACATT ATTCTCAGTG ATACAAGCGC CAGTGTCAGC
      SD      start relB
30     901  GAGCTGAAGA AAAACCCGAT GGCGACAGTC AGCGCCGGTG ATGGTTTCCC
      951  GGTGCTATC CTGAACCGTA ATCAGCCTGC TTTCTACTGT GTACCCGCAG
35    1001  AGCTGTACGA AAAGATGCTT GATGCCCTAG ACGATCAGGA GTTGTTTAAA
      1051  CTGGTAGCCG AACGCAGCAA CCAACCGCTG CATGATGTAG ATCTGGATAA
                                     SD
40     1101  end relB/start relE
      1151  ATACCTATGA GGTATCAGGT AAAATTCAGG GAAGATGCGC TGAAAGAGTG
      1201  GCAAAACTG GACAAGGCTA TTCAGCAACA GTTTGCGAAA AAGCTAAAAA
45     1251  AGTGCTGTGA CAATCCGCAT ATTCCTTCCG CAAAACTGCG TGGGATAAAG
      1301  GACTGCTACA AAATAAAATT ACGTGCCTCA GGTTCGCGC TGGTCTATCA
50     1351  GGTGATTGAC GAACAATTAA TTATCGCTGT TGTAGCTGTG GGTAACGTG
                                     end relE
      1351  AGCGCAGTGA CGTTTATAAT CTTGCCAGCG AAAGAATGAG ATAAAAGCAA

```

5

1401 TAAACACAGA AAGTTACTCT GCGTTATGG GGTAATGCAA AGTATGAGTC  
1451 GTAGAGGGAA TTGCCTGGAT AATTCGCCGA TGGAAAGAGT CTTTCGCAGC  
1501 CTTAAAAGTG AATGGCTTCC GAAAGGTGGT TATGGTGATT TTAGCCATGC  
(database accession number M26308)

### 1.3: Nucleotide sequence and proteins of a *relBE* homologous operon from *Bacillus thuringiensis*

Using BLAST database searching (Altschul et al., 1990) it was found that transposon

- 5 Tn5401 from the Gram-positive organism *B. thuringiensis* contains, in one end or asymmetrically located, a two-component system which exhibits both structural and sequence similarity with the above described *relBE* systems from *E. coli*. This homology is surprising given that it has not previously been described that *relBE*-like genes are found in organisms other than *E. coli*.

10

The nucleotide sequence of the *relBE* operon from Tn5401 is shown in Table 1.7. In this Table the transcriptional start-site of the *relBE* mRNA is indicated with an asterisk (Baum, 1994). IR indicates inverted repeats in the *relBE<sub>Bt</sub>* promoter region. Start codons and stop codons are shown in bold. The Shine & Dalgarno sequence of the *relB<sub>Bt</sub>* gene is doubly underlined. The DNA sequence is from Baum et al., 1989.

15

- The *relE<sub>Bt</sub>* gene codes for a very basic protein of 74 aa (pI = 10.6) and the *relB<sub>Bt</sub>* gene codes for an acidic protein of 87 aa (pI = 4.4). The protein sequences of RelE<sub>Bt</sub> and RelB<sub>Bt</sub> are aligned with the other RelE and RelB homologues in Tables 1.5 and 1.6, respectively. The modular, structural and physico-chemical similarities between the *B. thuringiensis* system and the *E. coli* systems suggested that the genes may exert similar functions in very different bacteria.

20

Table 1.7: DNA sequence of the *relBE* operon from the Gram-positive organism *B. thuringiensis* (SEQ ID NO:45)

```

5      3701  CTCGTTTTTT CTGTTGGTAC AAACCTAATT GATTTTGAAT AATTTGTTTG
      3751  TACCAGTCCT TTTTGCTTAG CCCAGTCAAA ATAACGTTTG ATTGAATTAA
      3801  TGCGCCGGTT AATCGTAGAA GGTTTTAGTA ATCTTGTAAC TTGCATATGC
10     3851  CCTCGATATC GAGCAATAGT GCGAGCGGTA ACTTCTATTG GATGAAAAAG
      3901  AGTATCCTCA GCATGTTTTT CCCACACATT TTCAAACCAA AATACAAAAT
      3951  CTTTTAAATC ACTCGTATAT TCTTTTAGTG TTTTGTATG CAAATCTCCT
15     4001  TCTTGAGATA AGCTAGAAAT AAAATCGGAA ATCAAAGATG TTGCTTGTAT
      4051  AGAAATTGTT TTAGTGGAAT GCATAAATAC CTCCTCTTTT ATTGACTTAC
                                     -35
20     4101  ATTAGCGGAC ATGATATTTT AATCTTATCA ATTATGTTAG CGGACATCAA
                                     -10      *→ relBEBt mRNA
      4151  ACATTTATTT TCCCACACTT CATGTCCACT AATATTAATT AGTGGACATT
                                     ----- → IR ← -----
                                     SD      Start relB
      4201  TAAACTATC TCGAAAGTAG GTGTAACACA TGGCTATTCTG TAAAGATGAA
30     4251  TTGTATCGGT TAATTGATCA CCTGGATCAA CAAGATGAAA AAGCAGCATT
      4301  TGACTTTTTA GAATTTCTTG TTCAACGGTC AAGAAGAAAA CCTAAAGAAT
35     4351  GGGAAAAAAT TGATATGGCA GATCCTGATC ATGAACCGCT GTCTACACAA
      4401  GAGTTAGAAC AGTTAAACAG TGAAGAAGGA TATGTATCAG GGGAGGACGC
      4451  AAAACGTGAA TTCGGACTAC AAATTGATTT ACCATAAGTC CGCGGTGAAA
                                     End relB      Start relE
45     4501  TTTATTGCAA AGCAAGAAAA AGGGATTCAA AAAAGAATTG CAGAAGGATT
      4551  GAAGGGACTT CTTAAGATTC CTCCTGAAGG AGATATTAAA AGTATGAAAG
50     4601  GTTACACAGA ACTATATCGA TTACGGATTG GAACCTTTCG AATTTTATTT

```

4651 GAAATAAATC ATGATGAGAA AGTCATATAC ATACAAGCAA TTGGAAATCG  
5                                      End *relE*  
4701 TGGTGACATC TATAAATAAG GCAAACATGC ATTTTAAAAA GAAAGGTCTT  
4751 CTGAATCGAA GAACCTTCCT TTTTGTGTG CGAATAATGT CCGCTAATGC  
10 4801 TTGTTGCGTG ATTCTGTTCC ATTGCTACAC ATACCCC  
      (database accession number U03554)

#### 1.4. The archaeon *Methanococcus jannaschii* encodes a *relBE* homologous system

Again using database searching it was found that the completely sequenced genome of the methanogenic archaeon *Methanococcus jannaschii* codes for three *relE* homologous genes, one of which are located just downstream of *relB* homologous genes. This finding was surprising since, in many respects, archaeal organisms are more similar to eukaryotes than to bacteria (e.g. in their macromolecular synthesis apparatuses).

- 10 The DNA sequence of the *relBE*<sub>Mj1</sub> system is shown in Table 1.8. In this Table start codons and stop codons are shown in bold. The DNA sequence is from Bult et al., 1996.

- Gene *relE*<sub>Mj1</sub> codes for a very basic polypeptide of 90 aa (pI = 11.0) and gene *relB*<sub>Mj1</sub> codes for an acidic polypeptide of 82 aa (pI = 4.4). The aa sequences of the RelE<sub>Mj1</sub> and RelB<sub>Mj1</sub> proteins are aligned with the other RelBE homologues in Tables 1.5 and 1.6, respectively. Thus, these basic similarities suggested that the *relBE*<sub>Mj1</sub> system may carry out similar or related functions in bacteria and archae. The properties of the second and third *relE* homologues of *M. jannaschii* are also given in Table 1.3. These comparisons show that *M. jannaschii* codes for one complete *relBE* homologous gene system and for two *relE* homologues without an adjacent *relB* partner.

Table 1.8. DNA sequence of a *relBE* homologous gene system from the archaeon*Methanococcus jannaschii* (SEQ ID NO:46)

5	751	CCGATACCGT	TGCTGGAGAC	ATAGCTGGAG	CTTTGAAGGC	GGAGAAGCTT
	801	ATTTTAATAA	CAGATGTTGA	TGGAATAATG	GATGATATAA	ATAATCCAGA
	851	GACGTTGCAT	AGAAAATTAA	CAGCTTCAGA	ACTAAAAGAA	ATGATAGAAG
10	901	ATGGAAGAAT	AAAGGGAGGG	ATGATTCCAA	AGGCTGAAAG	TGCCTTATAT
	951	GCCTTAGAGC	ATGGAGTTAA	GAGCGTTCAT	ATAATAAATG	GAAAGATTCC
15	1001	TCATGCTTTG	TTGTTGGAGA	TATTTACAGA	GGAGGGTATT	GGGACGATGA
	1051	TAACAAGAGA	TTAAAGTTTT	TATATTATAA	ACTACTTAAG	AATTAAAATA
				Start <i>relB<sub>Mj1</sub></i> :		
20	1101	AGACAAATAA	GGGGATAACT	<b>ATG</b> CTCAATA	TAAACAAAGA	GATAGCACAA
	1151	ATAGAACTG	AATTGAATGA	ATTGAAAAAA	TTGAGAGATG	AAATCTCTGA
	1201	AAGGATTGAA	AAATTAGAAA	TAAAGTTATT	AAAATTGAAA	GCATTAGCTA
25	1251	TTCCAGAGGA	GGAATTTGAA	GAGGATTATG	AAGAAATTAT	AGAAGATGTT
	1301	AAAAAATCTC	TGGATAAAAA	AGAGACTGTG	CCAGCAGAAG	AGGCTTTGAA
30			End <i>relB<sub>Mj1</sub></i> /start <i>relE<sub>Mj1</sub></i>			
	1351	AGAATTGGGA	TTATT <b>ATG</b> AA	GTTTAACGTT	GAGATACATA	AAAGAGTCTT
	1401	AAAAGATTTA	AAGGATTTGC	CTCCCTCAAA	CTTAAAGAAG	TTTAAAGAAC
35	1451	TAATAGAAAC	ATTAAAAACC	AATCCCATTG	CAAAAGAAAA	ATTTGATATT
	1501	AAAAGATTAA	AAGGCAGTGA	TGAGGTTTAT	AGAGTTAGAA	TTGGAAAATT
	1551	TAGAGTTCAA	TATGTTGTTT	TATGGGATGA	TAGAATAATA	ATAATTAGAA
40				End <i>relE<sub>Mj1</sub></i>		
	1601	AGATAAGTAG	AAGAGAAGGA	GCTTATAAAA	ATCCCT <b>TA</b> AGC	TATTAAAAAT
	1651	TCTAATGGCT	ACATTTTAT	ATCTCTTTTC	TTAATTCAAA	TAGAAAAAAC
45	1701	AGATTCGGCT	GATACCATGA	TTATTCTTTT	AGATTTAAAT	GGAACAATAG

(database accession number U67464)

### 1.5. *relBE* homologous genes are ubiquitous in prokaryotes

Further *relBE* homologous two-component systems were discovered. The corresponding RelB and RelE homologous proteins are aligned in Tables 1.5 and 1.6, respectively.

- 5 It appears that *relE* homologous genes are present in a wide variety of Gram-negative bacteria (*E. coli*, *H. influenzae*, *V. cholera*, *H. pylori* and *Synechosystis*), in Gram-positive bacteria (*B. thuringiensis* and *M. tuberculosis*) and in Archae (*M. jannaschii* and *A. fulgidus*). Most strikingly, the archaeon *A. fulgidus* contains four complete *relBE* homologous gene systems.

10

A number of features become evident from the alignments of the proteins (Tables 1.5 and 1.6) and from the properties listed in Tables 1.3 and 1.4. First, all RelE homologues are basic with pH's around 8-10 whereas the *RelB* homologues are acidic with pl's about 4-5. Secondly, the RelE proteins are in general slightly larger (90-120 aa)  
15 than the *RelB* homologues (70 - 80 aa). Thirdly, the start codons of the *relE* genes are juxtaposed or even overlap with the stop codons of the linked *relB* partner, thus indicating translational coupling of *relE* to *relB*. These properties suggest that the proteins could exert similar functions in very different organisms.

## 20 EXAMPLE 2

### Demonstration of translation of the *relB<sub>K-12</sub>* and *relE<sub>K-12</sub>* genes

- Using the low copy-number *lacZ* fusion vector pOU253 (Table 0.1) in frame gene fusions between *relB<sub>K-12</sub>* and *relE<sub>K-12</sub>* and the *lacZ* gene were constructed (see Materials  
25 and methods). Thus plasmid pKG4001 (+388 to +596) carries a fusion between *relB<sub>K-12</sub>* and *lacZ*, and pKG4002 (+388 to +921) carries a fusion between *relE<sub>K-12</sub>* and *lacZ*. The structure of the relevant parts of these reporter plasmids are shown in Fig. 1. When present in strain MC1000, both plasmids expressed significant amounts of -  
30 galactosidase fusion proteins, indicating that genes *relB<sub>K-12</sub>* and *relE<sub>K-12</sub>* are translated (Fig. 1). The *relE<sub>K-12</sub>-lacZ* fusion (pKG4002) expressed significantly lower amounts of  $\beta$ -galactosidase than the *relB<sub>K-12</sub>-lacZ* fusion, mainly because pKG4002 encodes an intact *relB<sub>K-12</sub>* gene which produces the RelB<sub>K-12</sub> autorepressor which inhibits transcription from the *relB* promoter.



## EXAMPLE 3

**Demonstration of translation of the *relB*<sub>P307</sub> and *relE*<sub>P307</sub> genes**

- 5 To detect authentic RelB<sub>P307</sub> and RelE<sub>P307</sub> proteins, *in vitro* translation reactions were carried out using high copy number pUC-plasmids carrying genes *relE*<sub>P307</sub> (pHA403), *relB*<sub>P307</sub> (pHA402) or both genes (pHA100) (for construction of these plasmids, see Materials and Methods and Table 0.1). Proteins produced in the *in vitro* translation reactions were labelled with <sup>35</sup>S-Methionine and separated by SDS-page. Fig. 2 shows
- 10 the direct visualization of RelB<sub>P307</sub> and RelE<sub>P307</sub>, thus providing evidence that the corresponding genes are translated.

## EXAMPLE 4

**15 Demonstrating that *relE*<sub>K-12</sub> is a cytotoxin**

- The low copy-number cloning vector pNDM220 contains *lacIq* and the LacI regulated pA1/O4/O3 promoter (Lanzer and Bujard, 1988) upstream of a multiple cloning site (mcs). The genetic structure of pNDM220 is shown in Fig. 3. Without IPTG added to
- 20 the growth medium, the pA1/O4/O3 promoter is almost completely turned off. However, with IPTG, strong transcription is induced towards the cloning site. Therefore plasmid pNDM220 is suitable for the conditional expression of genes, in particular toxin-encoding genes.
- 25 The *relE*<sub>K-12</sub> gene of *E. coli* K-12 (Bech et al., 1985) was PCR amplified and inserted into the mcs of pNDM220, resulting in pMG223 (for the construction of pMG223, see Materials and methods). Plasmid pMG223 (+733 to +1020) was established in MC1000, which contains a chromosomal copy of the *relBE* operon. However, it was not possible to transform pMG223 into the JS115 strain, which carries a deletion of
- 30 the *relBE* operon ( $\Delta relB$ ). Therefore, the induction experiments shown in Fig. 4 were accomplished using strain MC1000, which contains the chromosomal copy of *relBE*<sub>K-12</sub>.

Strain MC1000/pMG223 was grown in LB at 37°C. At time zero, IPTG was added to the growth-medium. After two hours of induction with IPTG, the viable counts decreased c. 600-fold (Fig. 4B). The decline started immediately and continued exponentially for about 2 hours. On plates containing IPTG, viable counts decreased even further (data not shown). The optical density (OD<sub>450</sub>) increased during the first 20 minutes after addition of IPTG and then the culture became stationary (Fig. 4A). Addition of IPTG to growing cells containing the vector-plasmid had no effect (not shown). These results indicate that the *relE* gene encodes a cell toxin.

## 10 EXAMPLE 5

### Demonstrating that RelB<sub>K-12</sub> is an antitoxin

Plasmid pMG2202 (+388 to +921) is a pBR322 derivative that contains the *relB* gene expressed from its own promoter (see Table 1.1). Plasmid pMG2201 (+388 to +597) is a pBR322 derivative that contains the *relB* promoter and the first part of *relB*<sub>K-12</sub>. Thus, pMG2201 does not contain an intact *relB* gene and was included in the analyses as a control plasmid. The strains MC1000/pMG223 (pA1/O4/O3::*relE*+) /pMG2202 (*relB*<sup>+</sup>) and MC1000/pMG223/pMG2201 (*relB*) were subjected to a physiological growth experiment similar to the one described in Example 4. As seen from Fig. 4E and 4F, the presence of the high copy-number *relB*-carrying plasmid suppressed *relE*-dependent cell killing. The antitoxin effect was dependent on an intact *relB* reading frame, since the control-plasmid (pMG2201) carrying the promoter region and the first part of the *relB* reading frame did not prevent the *relE* mediated cell killing (Fig. 4C, 4D).

## EXAMPLE 6

### Demonstrating that *relE*<sub>P307</sub> encodes a very efficient cytotoxin

30

The medium copy number expression vector pBAD33 contains an arabinose inducible promoter (pBAD) with a multiple cloning site (mcs) and the *araC* gene (Guzman et al., 1995). The genetic structure of pBAD33 is shown in Fig. 5. Without arabinose added to the growth medium, the pBAD promoter is completely turned off. However, with

arabinose, strong transcription is induced towards the cloning site. On top of this property, the pBAD promoter is repressible by the addition of glucose to the growth medium. Thus, by the addition of glucose, transcription from pBAD can be rapidly and efficiently turned off.

5

The glucose repression effect is epistatic to the inducer effect by arabinose. Hence, if cells with a pBAD-carrying plasmid are grown in a medium containing both arabinose and glucose then the promoter is not induced. However, if cell-growth depletes the medium for glucose, then the promoter will be induced. Therefore, plasmid pBAD33 is  
10 suitable for the conditional turning on and off of the expression of genes, in particular toxin-encoding genes as described herein.

The *relE* gene of the *E. coli* plasmid P307 (Saul et al., 1989) was PCR amplified and inserted into the mcs of pBAD33, resulting in pHA810 (for the construction of plasmid  
15 pHA810, see Materials and methods). Thus plasmid pHA810 contains the *relE*<sub>P307</sub> gene inserted downstream of the pBAD promoter. Strain MC1000/pHA810 was grown in LB-medium without glucose at 37°C. At time zero, the culture was diluted into medium containing either 0 or 0.2% arabinose. In the arabinose-containing culture, an immediate decline in viable counts was observed (Fig. 6B, closed symbols).  
20 The decline continued exponentially throughout the experiment. After 240 min of induction with arabinose, viable counts had decreased more than five orders of magnitude. Without arabinose, cells containing pHA810 continued to grow exponentially (Fig. 6A and 6B, open symbols). On plates containing arabinose, none or very few viable cells were detected. These results show that *relE*<sub>P307</sub> gene encodes an extremely  
25 efficient cell toxin.

#### EXAMPLE 7

##### Demonstrating that RelB<sub>P307</sub> is an antitoxin

30

Plasmid pHA110 (+1 to +1122) is a pBR322 derivative that contains the *relB*<sub>P307</sub> gene expressed from its own promoter. The strain MC1000/pHA810/pHA110 (*relB*<sub>P307</sub>) was subjected to a physiological growth experiment as described in Example 6. It appeared that the presence of the *relB*<sub>P307</sub>-carrying plasmid pHA110 prevented

*relE*<sub>P307</sub> dependent inhibition of cell growth (Fig. 6C) and cell killing (Fig. 6D). This observation shows that *relB*<sub>P307</sub> codes for an antitoxin that counteracts the cell killing caused by RelE<sub>P307</sub>.

## 5 EXAMPLE 8

### Determination of the frequency of spontaneous mutants that are resistant to the killing effect of RelE

- 10 Strain MC1000/pHA810 was grown exponentially to an OD of 0.5 and serial dilutions of the cell suspension were plated on LA plates containing chloramphenicol (selecting for plasmid pHA810) and with or without 0.02% arabinose (which induces expression of *relE* present in pAH810). On such plates without arabinose the plating efficiency of strain MC1000/pAH810 was normal, i.e. more than 99% of the viable cells produced
- 15 a colony. This indicated that the presence of pAH810 in itself had no effect on the viability of the cells. However, with arabinose the plating efficiency was reduced by about 10<sup>9</sup> fold, thus indicating that expression of RelE is extremely toxic to the cells. The few surviving colonies that appeared eventually were retransformed with the RelE expression plasmid pHA210 which can co-exist with pAH810. However, none of the
- 20 surviving cells from the first round of selection (i.e. using pHA810) survived induction of RelE (by addition of IPTG) from the second plasmid pAH210.

These results show that resistance against RelE toxicity is a very rare event, as based on this experiment it is less than about 10<sup>-9</sup>.

25

## EXAMPLE 9

### Demonstrating that RelE of the Archeon *Methanoccus jannaschii* is toxic to *E. coli*.

- 30 The *relE* gene of *M. jannaschii* was amplified from genomic DNA using primers MJ-relE/2CWW (5'-CCCCCGAATTCGCATGCGCCATTAGAAT, SEQ ID NO:47) and MJ-relE/1CW (5'-CCCCCGGATCCGAGCTCGAGGCTTTGAAAGAATTGGG, SEQ ID NO:48). The resulting DNA fragment was cleaved with *Bam*HI and *Eco*RI and cloned into plasmid pNDM220 (Fig. 3) thus yielding pHA705 (Fig. 7). Similarly, *relB* and *relE*

from *M. jannashii* were PCR amplified using primers relB-M.jannCW (5'-CCCCGGATC-CGTCGACGACAAATAAGGGGATAACTATG, SEQ ID NO:49) and MJ-relE/2CWW. The resulting DNA fragment was cleaved with *Bam*HI and *Eco*RI and cloned into pNDM220, thus yielding pHA715 (Fig. 7).

5

Plasmids pHA705 (carrying *relE*) and pHA715 (carrying *relBE*) were transformed into *E. coli* K-12 strain MC1000. Cells were grown exponentially and followed after the addition of IPTG. Fig. 8 shows that the addition of IPTG inhibited the growth of MC1000/pHA705 but not that of MC1000/pHA715, and Fig. 9 shows that viable  
10 count was significantly reduced in the case of MC1000/pHA705 but not in that of MC1000/pHA715, thus demonstrating that RelE of *M. jannashii* is toxic to *E. coli*.

#### EXAMPLE 10

15 **Demonstrating that RelE of the Gram-positive bacterium *Streptococcus pneumoniae* is toxic to *E. coli*.**

Using BLAST database searching, we identified two homologues of the *relBE* genes of *S. pneumoniae*. The DNA sequence of the homologue designated *relE<sub>Sp2</sub>* is shown in  
20 Fig. 10. Gene *relE<sub>Sp2</sub>* was PCR amplified from genomic DNA of *S. pneumoniae* strain RP46 using primers relE-Sp2/cw (5'-CCCCGGATCCGATGCATGATTAGGCTTGAAG, SEQ ID NO:50) and relE-Sp2/ccw (5'-CCCCGAATTCGAATGAAA-  
ATTACTTGAAAAAAG, SEQ ID NO:51). The resulting DNA fragment was cleaved with *Bam*HI and *Eco*RI and cloned into pNDM220 thus yielding plasmid pHA-Sp2 (Fig.  
25 11).

Plasmid pHA-Sp2 (carrying *relE<sub>Sp2</sub>*) was transformed into *E. coli* strain MC1000. Cells were grown exponentially and followed after the addition of IPTG. Fig. 8 shows that the addition of IPTG inhibited the growth of MC1000/pHA-Sp2, and Fig. 9 shows that  
30 viable counts were dramatically reduced, thus demonstrating that expression of *relE<sub>Sp2</sub>* is highly toxic to *E. coli*.

## EXAMPLE 11

**Cloning of the *relE* genes of plasmid P307, *M. jannashii* and *E. coli* K-12 into the broad-host-range vector pHAG33.**

5

The broad-host-range vector pVLT33 is an RSF1010 derivative that can be mobilized by an appropriate conjugation system (de Lorenzo, Eltis, L., Kessler, B. and Timmis, K.N. 1993. Analysis of *Pseudomonas* gene products using *lacI<sup>P</sup>/Ptrp-lac* plasmids and transposons that confer conditional phenotypes. Gene 123, 17-24). It also contains  
10 the *tac*-promoter (*ptac*) and *lacI<sup>P</sup>*. Since *ptac* is leaky and therefore unsuitable for the regulated expression of toxins, the promoter was replaced by the pA1/O4-O3 promoter of pNDM220. The resulting plasmid, pHAG33, is shown in Fig. 12. The *relE* genes of pHA210 (*relE<sub>P307</sub>*), pHA705 (*relE<sub>Mj</sub>*) and pMG223 (*relE<sub>K12</sub>*) were cloned into pHAG33, resulting in plasmids pHA33-2 (Fig. 13), pHA33-3 (Fig. 14), and pHA33-4  
15 (Fig. 15), respectively.

## EXAMPLE 12

**Demonstrating that RelEs of *E. coli* K-12, P307 and *M. jannashii* are toxic to *Pseudomonas putida*.**

20

Plasmids pHA33-2 (*relE<sub>P307</sub>*), pHA33-3 (*relE<sub>Mj</sub>*) and pHA33-4 (*relE<sub>K12</sub>*) were transformed into the *E. coli* K-12 strain S17-1. This strain contains the conjugation system of RP4 and is thus able to mobilize pHA33-derived plasmids as described above (Simon et al.,  
25 1986). After conjugation on solid medium to *P. putida* strain KT2440 according to standard procedure, strains KT2440/pHA33-2, KT2440/pHA33-3 and KT2440/pHA33-4 were established.

The strains were grown exponentially in LB containing 30 µg/ml ampicillin and 50  
30 µg/ml kanamycin and followed after the addition of 2 mM IPTG. As seen from Fig. 16, the increment in cell-growth as measured by OD<sub>450</sub> was reduced by IPTG in all three cases. Furthermore, measurements of viable counts (Fig. 17) showed cell-killing in all three cases, most severe in the case of *relE<sub>K12</sub>* (pHA33-4(+)) in Fig. 17). Thus, RelE

proteins of P307 and *M. jannashii* are toxic to *P. putida* and RelE of *E. coli* K-12 is extremely toxic to *P. putida*.

### EXAMPLE 13

5

#### Demonstrating biological containment by the depletion of a carbon source.

Plasmid pHA810 (Fig. 18) was constructed by inserting the *relE* gene of P307 into the expression vector pBAD33 (Fig. 5). The promoter (designated  $p_{BAD}$ ) upstream of  
10 *relE*<sub>P307</sub> in pHA810 is repressed by glucose and induced by arabinose. The repression by glucose overrides induction by arabinose such that the simultaneous presence of glucose and arabinose in the growth medium results in repression of the promoter.

To simulate a realistic scenario in which the carbon source was depleted, we grew  
15 MC1000/pHA810 in ABT minimal salts medium at 35°C in the presence of a limiting amount of glucose (0.025% w/v) (represses  $p_{BAD}$ ) and varying the concentration of arabinose (induces  $p_{BAD}$ ). Optical density (Fig. 19) and viable counts (Fig. 20) typical for such an experiment were obtained. As seen in Fig. 19, the rate of increase in OD<sub>450</sub> is severely reduced by the highest amounts of arabinose (0.050% and  
20 0.075%). This was expected, since arabinose induces  $p_{BAD}$  and the limited amount of glucose (0.025%) cannot fully suppress  $p_{BAD}$  at high concentrations of arabinose. The glucose added (0.025%) was depleted by cell growth at an OD<sub>450</sub> = approx. 0.1. At this OD<sub>450</sub>, a dramatic cell killing was seen in the case of 0.005%, 0.010%, and 0.025% of arabinose (Fig. 20). This result shows, that depletion of the carbon source  
25 (glucose) leads to massive cell killing, and thus to biological containment of the plasmid that carries *relE*<sub>P307</sub>.

### EXAMPLE 14

#### 30 Demonstrating that RelE of *E. coli* K-12 and *M. jannashii* are toxic to human cells.

The cell line 293 is a permanent line of primary human embryonal kidney cells transformed by human adenovirus type 5 (Ad 5) DNA (ATCC CRL-1573). The cells are particularly sensitive to human adenovirus, are highly permissive for adenovirus DNA, and

contain and express the transforming genes of AD 5 (Graham, F.L., Smiley, J., Russell, W.C., and Nairn, R. 1977. Characteristics of a human cell line transformed by DNA from human adenovirus type 5. J. Gen. Virol. 36, 59-74).

- 5 Plasmid pcDNA3.1(+) (Invitrogen) carries the constitutive promoter  $P_{CMV}$  from cytomegalovirus upstream of a multiple cloning site (mcs). Genes *relE* of *E. coli* K-12 and *M. jannashii* were inserted in the mcs, resulting in plasmids p5.4 and p5.3, respectively.
- 10 Plasmids pcDNA3.1(+)(control), p5.4 and p5.3 were transfected into cell line 293 by selection in medium containing G418 (geneticin), which selects for cells expressing the neomycin gene present on the plasmids. After 12 days, the cell density was measured by inspection. In the case of p5.4 (*relE*<sub>K-12</sub>), between 0 and 5% of the cells had survived (as compared to the control). In the case of p5.3 (*relE*<sub>Mj</sub>), between 5 and 15 10% of the cells had survived. These results indicate that the bacterial RelE<sub>K-12</sub> and the archaeal RelE<sub>Mj</sub> toxins both are lethal to human cells.

#### EXAMPLE 15

#### 20 Demonstrating that RelE<sub>K-12</sub>, RelE<sub>P307</sub>, and RelE<sub>Mj</sub> inhibit translation in vitro

DNA fragments comprising genes *relE*<sub>K-12</sub>, *relE*<sub>P307</sub> and *relE*<sub>Mj</sub> were PCR amplified such that a T7 RNA polymerase promoter was placed upstream of the corresponding genes (according to Thisted et al., 1994).

25

- The following primers were used: *relE*<sub>K-12</sub> (P1: 5'-TGTAATACGACTCACTATAGATAAGGAGTTTTATAAATGGCGTATTTCTGGATTTTG, SEQ ID NO:52) and P2 (CACCTTCGGTGCGAAACAG, SEQ ID NO:53); *relE*<sub>P307</sub> (P3: 5'-TGTAATACGACTCATATAGATAAGGAGTTTTATAAATGAGGTATCAGGTAAAATTCA (SEQ ID NO:54) and 30 P4: 5'-CTTTCATCGGCGAATTATC, SEQ ID NO:55); *relE*<sub>Mj</sub> (P5: 5'-TGTAATACGACTCACTATAGATAAGGAGTTTTATAAATGAAGTTTAACGTTGAGATAC SEQ ID NO:56) and P6: (5'-ATCATGGTATCAGCCGAATC, SEQ ID NO:57). T7 RNA polymerase sequences are underlined, and the strong Shine-Dalgarno (SD) sequence from the *parA* system of plasmid R1 is shown in italics.



Using in vitro transcription with T7 RNA polymerase according to standard procedures, mRNAs encoding *relE*<sub>K-12</sub>, *relE*<sub>P307</sub>, and *relE*<sub>Mj</sub> were produced and subsequently purified from a denaturing polyacrylamide gel. To facilitate the quantification of the mRNAs they were labelled with tritium (alpha-<sup>3</sup>H-CTP) during their synthesis. The *relE*-  
5 encoding mRNAs (1.5 pmol) were used as templates in in vitro translation reactions employing an S30 extract (obtained from Promega) containing 150 µM of each amino acid except Methionine which was 1 µM.

Figure 21 shows SDS-PAGE (tricine-gel) analysis of such an experiment. The in vitro  
10 translation reactions were initiated with unlabelled Methionine in order to produce RelE toxin in the reaction. Ten minutes after the addition of the unlabelled Methionine, radioactive 35-S-Methionine (5 pmol in a 15 µl volume) was added and the reaction continued for an additional 20 minutes. C in Fig. 21 denotes a control lane without exogenous mRNA added. The protein bands seen in this lane originate from translation  
15 of mRNAs present in the S30 extract. In lane 1, the *in vitro* translation reaction contained an mRNA encoding the *relE* gene of *E. coli* K-12. As seen, the translation reaction was severely inhibited. In lane 2, a mRNA encoding a mutated *relE* (denoted *relEmE* and described in Gotfredsen et al, 1998) gene was added. As seen, the presence of this mRNA did not inhibit the reaction. This result shows that the RelE protein  
20 produced during the initial incubation-period without 35-S-Met added inhibits the in vitro translation reaction (i.e. compare lanes 1 & 2). Furthermore, this lack of inhibition is correlated with loss of cell killing activity in vivo (since the mutated *relE* gene, *relEmE*, used in lane 2 is not toxic to *E. coli* cells), thus indicating that inhibition of translation is the actual cause of cell death in vivo. In lanes 3 and 4, mRNAs encoding  
25 *relE* of plasmid P307 and the archaeon *M. jannashii* were added. As seen, the presence of these mRNAs inhibited the in vitro translation reactions as well. These results indicate that the RelE toxins from *E. coli* K-12, *M. jannashii* and plasmid P307 all act by inhibition of translation.

## EXAMPLE 16

Demonstrating that RelE<sub>K-12</sub> is toxic to yeast cells.

5 1. Yeast strain

In these experiments the yeast strain *Saccharomyces cerevisiae* 281288DIV-36 (MATa his 4-5; LEU2 THR4 ura3-52 trp1 CYH2 KAR1) was used.

10 2. PCR amplification of RelE coding region

The RelE coding region was PCR amplified from the plasmid pMG223 using two oligonucleotide primers. The primer S-RelE was 24 nucleotides long (5'-TAGGTACCATGGCGTATTTTCTGG-3', SEQ ID NO:58). It contains *KpnI* and *NcoI* endonuclease restriction sites at the 5' end with an 8 nucleotide overhang. Primer AS-RelE was 23 nucleotides long (5'-GAGACCCCACACTACCATCGGCG-3', SEQ ID NO:59) and hybridises 400 nucleotides downstream the RelE termination codon and 392 nucleotides downstream the *EcoRI* site in plasmid vector pMG223. PCR amplifications were performed using Vent® Polymerase (New England Biolabs), 200 μM of each dNTP. PCR reaction buffer (10 mM KCl; 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>; 20 mM Tris-HCl (pH 8.8); 2 mM MgSO<sub>4</sub>; 0.1% Triton X-100) with 0.2 μM of each of the primers. After 5 min denaturation (95°C) PCR was performed with 20 cycles, each cycle consisting of 1 min denaturation (92°C), 1 min primer annealing (50°C) and 1 min primer extension (72°C). Successful PCR products of 706 bp DNA fragments were identified and purified from a 1% agarose gel after a run of 1 hour at 40 mA. The PCR product was digested with the two restriction enzymes *KpnI* and *EcoRI* and the fragment of 304 bp containing the RelE open reading frame was isolated after electrophoresis on a 1.2 % agarose gel, and purified using a gel extraction kit (Pharmacia).

### 3. Cloning of the amplified *relE* gene

The isolated DNA fragment of the *relE* gene flanked with *KpnI* and *EcoRI* sites was ligated into the pYES2 expression vector (Invitrogen) previously digested with *KpnI* and *EcoRI* using standard procedures (Sambrook). After ligation, *E. coli* Top10 (Invitrogen) was transformed with the ligation mixture using electroporation using the *E. coli* gene pulser (BioRad). After phenotypic expression for 2 hours in SOC medium the culture was spread onto selective LB medium (Sambrook) containing 50 µg of ampicillin per ml. Transformed colonies were identified using PCR amplification and a positive clone designated pPK727 was further tested by restriction enzyme analysis. The functionality of the PCR amplified *RelE* gene was tested in *E. coli* by cloning the *NcoI-EcoRI* fragment from pPK727 into the *E. coli* expression vector pUHE24. Induction with IPTG led to cell killing in *E. coli*.

### 4. Yeast transformation

*S. cerevisiae* was grown overnight (ON) in YDP medium (1% yeast extract; 2% Bacto peptone; 2% glucose). For a single transformation, cells from 1 ml ON culture were spun down (5,000 rpm for 30 sec using an Eppendorf minicentrifuge) and washed twice in sterile water. Cells were resuspended in 200 µl lithium acetate buffer (10 mM Tris-HCl pH 7.6 with 100 mM LiOAc, 1 mM EDTA). After incubation for 15 min. at 25°C with agitation two transformations were made adding 20 µl carrier DNA (10 mg/ml salmon sperm DNA, sonicated and heat denatured) and 100 ng of the plasmids pPK727 and pYES2 (vector control), respectively. A volume of 1.2 ml 40% PEG 4,000 in 0.1 M lithium acetate buffer was added to each transformation mixture. The transformation mixtures were incubated in a 25°C incubator for 30 min before transferring to a 42°C water bath for 15 min. Cells were spun down (5,000 rpm for 30 sec.) and washed once with sterile water before plating on Uracil drop-out medium (1% Bernstein acid; 0.1% NaOH; 2% glucose; 0.67% Bacto yeast nitrogen base; 0.1% amino acids (without uracil); 2% agar).

After three days growth at 30°C single colonies were picked and streaked onto plates with Uracil drop-out medium. After two days at 30°C cells were transferred to induc-

tion medium (Uracil drop-out medium with 2% galactose as the sole carbon source) by replica-plating.

Single colonies of *S. cerevisiae* containing pPK727 and pYES2 were transferred to liquid Uracil drop-out medium (1% Bernstein acid; 0.1% NaOH; 2% glucose; 0.67% Bacto-yeast nitrogen base; 0.1% amino acids (without uracil)) and incubated ON. To compensate for difference in cell density, a volume of 50  $\mu$ l per OD<sub>540</sub> (optical density at 540 nm) was used to inoculate 50 ml of liquid Uracil drop-out medium with either glucose or galactose as sole carbon source, respectively. The four flasks with *S. cerevisiae* (pPK727) and *S. cerevisiae* (pYES2) in Uracil drop-out medium with or without galactose were incubated at 30°C with moderate shaking (200 rpm). To monitor growth, samples were taken at different time point and OD<sub>540</sub> was determined. Samples were taken in duplicates and the average OD<sub>540</sub> calculated and plotted against time of sampling

15

## 5. Results.

All colonies containing either the plasmid pPK727 or the pYES2 control plasmid were able to grow on plates with glucose as carbon source. When transferred to plates with galactose as the sole carbon source leading to gene expression from the P-gal1 promoter only cells with the pYES2 control plasmid showed normal growth, whereas cells containing the pPK727 were strongly inhibited in growth.

In liquid media, the yeast cells in which the *relE* gene was induced showed a remarked growth inhibition when compared to the uninduced control and to the controls with only the plasmid pYES2 without insert.

These results that are summarised in the below Table 16.1 clearly suggest that inhibition of cell growth in yeast cells is due to expression of the *relE* gene.

30

Table 16.1. Growth of *Saccharomyces cerevisiae* transformed with pYES2 +/- *relE* gene

time (hours)	Plasmid + relE		plasmid (control)	
	Galactose	no galactose	galactose	no galactose
0	0	0	0	0
5,5	0,018	0,024	0,023	0,028
15,5	0,013	0,186	0,055	0,176
23	0,190	1,195	0,286	0,963
28	0,010	1,480	0,816	1,649
29	0,035	3,930	1,660	2,650
65	0,990	3,950	6,180	3,820

## REFERENCES

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment search tool. *J Mol Biol* **215**:403-410.
- 5 Baum J.A. 1994. Tn5401, a new class II transposable element from *Bacillus thuringiensis*. *J Bacteriol* **176**:2835-2845.
- Bech F.W, Jorgensen S.T, Diderichsen B, Karlstrom O.H. 1985. Sequence of the *relB*  
10 transcription unit from *Escherichia coli* and identification of the *relB* gene. *EMBO J* **4**:1059-1066.
- Bertani, G. (1951). The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol* **62**:293-300.
- 15 Bolivar, F. (1978). Construction and characterization of new cloning vehicles. III. Derivatives of plasmid pBR322 carrying unique *EcoRI* sites for selection of *EcoRI* generated recombinant DNA molecules. *Gene* **4**:121-136.
- 20 Bult C.J, White O, Olsen G.J, Zhou L, Fleischmann R.D, Sutton G.G, Blake J.A, Fitzgerald L.M, Clayton R.A, Gocayne J.D, Kerlavage A.R, Dougherty B.A, Tomb J.F, Adams M.D, Reich C.I, Overbeek R, Kirkness E.F, Weinstock K.G, Merrick J.M, Glodek A, Scott J.L, Geoghagen N.S.M, Weidman J.F, Fuhrmann J.L, Venter J.C et al. 1996. Complete genome sequence of the methanogenic archaeon, *Methanococcus* jan-  
25 *naschii*. *Science* **273**:1058-1073.
- Casadaban M.J, Cohen S.N. 1980. Analysis of gene control signals by DNA fusion and cloning in *Escherichia coli*. *J Mol Biol* **138**:179-207.
- 30 Clark, D. and Maaløe, O. 1967. DNA replication and the division cycle in *Escherichia coli*. *J Mol Biol* **23**:99-112.

- Dam, M. and Gerdes, K. 1994. Partitioning of plasmid R1: Ten direct repeats flanking the *parA* promoter constitute a centromere-like partition site *parC*, that expresses incompatibility. *J. Mol. Biol.* **236**:1289-1298.
- 5 Fleischmann R.D, Adams M.D, White O, Clayton R.A, Kirkness E.F, Kerlavage A.R, Bult C.J, Tomb J.F, Dougherty B.A, Merrick J.M et al. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496-512.
- Gerdes, K., Bech, F.W., Jørgensen, S.T., Løbner-Olesen, A., Atlung, T., Boe, L., Karlstrøm, O., Molin, S. and von Meyenburg, K. 1986. Mechanism of postsegregational killing by the *hok* gene product of the *parB* system of plasmid R1 and its homology with the *relF* gene product of the *E. coli relB* operon. *EMBO J.* **5**:2023-2029.
- 10 Gotfredsen, M and Gerdes, K. 1998. The *Escherichia coli relBE* genes belong to a new toxin-antitoxin gene family. *Molec. Microbiol.*, **29**:1065-1076.
- Guzman L.M, Belin D, Carson M.J, Beckwith J. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J Bacteriol* **177**:4121-4130.
- 20 Jensen, R.B. and Gerdes, K. 1995. Microreview. Programmed cell death in bacteria: proteic plasmid stabilization systems. *Mol Microbiol* **17**:205-210.
- Klenk H.P, Clayton R.A, Tomb J.F, White O, Nelson K.E, Ketchum K.A, Dodson R.J, Gwinn M, Hickey E.K, Peterson J.D, Richardson D.L, Kerlavage A.R, Graham D.E, Kyrpides N.C,
- 25 Fleischmann R.D, Quackenbush J, Lee N.H, Sutton G.G, Gill S, Kirkness E.F, Dougherty B.A, McKenney K, Adams M.D, Loftus B, Venter J.C et al. 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* **390**:364-370.
- 30 Lanzer M, Bujard H. 1988. Promoters largely determine the efficiency of repressor action. *Proc Natl Acad Sci U.S.A* **85**:8973-8977.

Miller, J.F. (1972). Experiments in molecular genetics. Cold Spring Harbor Laboratory Press, New York.

Minton N.P. 1984. Improved plasmid vectors for the isolation of translational lac gene  
5 fusions. *Gene* **31**:269-273.

Saul D, Spiers A.J, McAnulty J, Gibbs M.G, Bergquist P.L, Hill D.F. 1989. Nucleotide  
sequence and replication characteristics of RepFIB, a basic replicon of IncF plasmids. *J*  
*Bacteriol* **171**:2697-2707.

10

Shägger, H. and von Jagow, G. (1987). Tricine-Sodium Dodecyl Sulfate-polyacryla-  
mide gel electrophoresis for the separation of proteins in the range from 1 to 100 kD.  
*Anal Biochem* **166**:368-379.

15 Simon, R., O'Connell, M., Labes, M. and Puhler, A. 1986. Plasmid vectors for the ge-  
netic manipulation of rhizobia and other gram-negative bacteria. *Methods. Enzymol.*  
**118**:640-659.

Thisted, T., Nielsen, A. and Gerdes, K. 1994. Mechanism of post-segregational killing:  
20 translation of Hok, SrnB, and PndA mRNAs of plasmids R1, F and R483 is activated  
by 3'-end processing. *EMBO J.* **13**:1950-1959.



## CLAIMS

1. A method of conditionally controlling the survivability of a recombinant microbial cell population, the method comprising (i) providing in the cells of said population a  
5 gene coding for a cytotoxic first kind of polypeptide, the gene is selected from the group consisting of the gene coding for the *E. coli* K-12 RelE polypeptide and a gene coding for a functionally equivalent polypeptide (said genes collectively being designated herein as the *relE* gene family), said gene is expressible in the cells of the population and, operably linked to the gene, a regulatable regulatory DNA sequence and (ii)  
10 cultivating the cell population under conditions where the *relE* gene or the gene coding for a functionally equivalent polypeptide is expressed, the expression leading to an at least partial killing of the cell population.
2. A method according to claim 1 wherein the gene coding for the functionally  
15 equivalent polypeptide is derived from a Gram-negative bacterium.
3. A method according to claim 2 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-negative bacterium selected from the group consisting of *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp.,  
20 *Pseudomonadaceae* spp., *Helicobacter* spp. and *Synechosystis* spp.
4. A method according to claim 1 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-positive bacterium.
- 25 5. A method according to claim 4 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-positive bacterium selected from the group consisting of lactic acid bacterial spp., *Bacillaceae* spp. and *Mycobacterium* spp.
6. A method according to claim 5 wherein the Gram-positive bacterium is *Bacillus*  
30 *thuringiensis*.
7. A method according to claim 1 wherein the gene coding for the functionally equivalent polypeptide is derived from a species belonging to *Arhae*.

8. A method according to claim 1 wherein the *relE* family gene is present on the chromosome of the cells.
9. A method according to claim 8 wherein the cells of the population do not contain a gene coding for a second type of polypeptide that is capable of counteracting the cell toxic effect of the RelE polypeptide or the functional equivalent hereof.
10. A method according to claim 1 wherein the gene coding for the RelE polypeptide or the functional equivalent hereof is present on an extrachromosomal replicon.
- 10 11. A method according to claim 1 wherein the regulatory DNA sequence regulates the expression of the gene coding for the RelE polypeptide or the functionally equivalent polypeptide at the transcriptional level by means of a promoter, the function of which is regulated by the presence or absence of a chemical compound in the cultivation medium.
- 15 12. A method according to claim 11 wherein the promoter is inducible by a chemical compound.
- 20 13. A method according to claim 11 wherein the promoter is suppressible by a first kind of chemical compound and inducible by a second kind of chemical compound whereby, when the first kind of compound is depleted from the medium, the promoter is induced by the second kind of compound.
- 25 14. A method according to claim 1 wherein the cells of the population comprises a gene coding for a second kind of polypeptide that is capable of binding to the *relE* polypeptide or the functional equivalent, said binding resulting in that the toxic effect of the RelE polypeptide or the functional equivalent is at least partially counteracted.
- 30 15. A method according to claim 14 wherein the gene coding for the second kind of polypeptide is operably linked to a regulatable regulatory DNA sequence, permitting that the gene coding for the second kind of polypeptide is suppressed under conditions where the gene coding for the RelE polypeptide or the functional equivalent is expressed.

16. A method according to claim 14 wherein the second kind of polypeptide is the RelB polypeptide derived from *E. coli* K-12 or a functionally equivalent polypeptide.

17. A method according to claim 16 wherein the functionally equivalent second kind of polypeptide is derived from the group consisting of a Gram-positive bacterium, a Gram-negative bacterium and *Archae* spp.

18. A method according to claim 1 wherein the expression of the *relE* gene family gene is stochastically regulated.

10

19. A method according to claim 18 wherein the stochastic regulation is effected by operably linking the *relE* gene family gene to a regulatory sequence that comprises an invertible promoter.

15 20. A method according to claim 18 wherein the stochastic regulation is effected by flanking at least part of the regulatory sequence by repeat sequences whereby the at least part of the regulatory sequence is recombinationally excised.

21. A method according to claim 1 wherein the cells contain a gene coding for a gene product of interest.

20

22. A method according to claim 21 wherein the gene product of interest is an immunologically active gene product.

25 23. A method according to claim 21 wherein the gene product of interest is one that is effective in degradation of an environmental pollutant.

24. A method according to claim 21 wherein the gene product of interest is a pesticidally active product.

30

25. A method according to claim 24 wherein the gene coding for the pesticidally active gene product is derived from *Bacillus thuringiensis*.

26. A method of confining an extrachromosomal replicon to a microbial cell population, the method comprising the steps of

(i) isolating a microbial cell naturally containing a gene belonging to the *relE* gene family coding for a first kind of polypeptide that, when it is expressed in the cell, acts as a toxin for the cell or, if the cell does not naturally contain a gene belonging to the *relE* gene family, introducing such a gene into the cell,

(ii) introducing into the cell the extrachromosomal replicon to be confined, said replicon containing a gene coding for a second kind of polypeptide that, by binding to the first kind of polypeptide, acts as an antitoxin for said first kind of polypeptide,

(iii) cultivating the cell under conditions where the genes coding for the first and the second kind of polypeptides are expressed, whereby a daughter cell that does not receive a copy of the extrachromosomal replicon is killed by the first kind of polypeptide being expressed in the absence of expression of the second kind of polypeptide.

27. A method according to claim 26 wherein the cell population consists of cells that comprises a gene coding for a gene product of interest.

28. A method according to claim 27 wherein the gene product of interest is selected from the group consisting of an enzyme, an immunologically active polypeptide, a pesticidally active gene product and a pharmaceutically active gene product.

29. A method according to claim 26 wherein the replicon is a plasmid occurring in the microbial cells at a copy number which is in the range of 1-30 including the range of 1-10 such as the range of 1-5.

30. A method according to claim 26 wherein the microbial cells belong to a Gram-negative bacterial species.

31. A method according to claim 30 wherein the bacterial species is selected from the group consisting of *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp. and *Pseudomonadaceae* spp.

32. A method according to claim 26 wherein the microbial cells belong to a Gram-positive bacterial species.

33. A method of post-segregationally stabilizing a plasmid in a microbial host cell  
5 population, the method comprising the steps of

(i) recombinationally inserting into the plasmid (a) a gene coding for a first kind of polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide and a functional equivalent thereof, said first kind of polypeptide having a toxin effect  
10 on the host cell and (b) a gene coding for a second kind of polypeptide that (1) is capable of acting as an antitoxin for first kind of polypeptide and (2) is capable of being degraded in the host cell at a higher rate than that at which the first kind of polypeptide is degraded,

15 (ii) cultivating the cell population under conditions where the genes coding for the first kind and second kind of polypeptides are expressed,

whereby a daughter cell that does not receive at least one copy of the plasmid is killed as a result of the faster degradation of the second kind of polypeptide.

20

34. A method according to claim 33 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-negative bacterium.

35. A method according to claim 34 wherein the gene coding for the functionally  
25 equivalent polypeptide is derived from a Gram-negative bacterium selected from the group consisting of *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp., *Pseudomonadaceae* spp., *Helicobacter* spp. and *Synechosystis* spp.

36. A method according to claim 33 wherein the gene coding for the functionally  
30 equivalent polypeptide is derived from a Gram-positive bacterium.

37. A method according to claim 36 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-positive bacterium selected from the group consisting of lactic acid bacterial spp., *Bacillaceae* spp. and *Mycobacterium* spp.

38. A method according to claim 37 wherein the Gram-positive bacterium is *Bacillus thuringiensis*.

39. A method according to claim 33 wherein the gene coding for the functionally  
5 equivalent polypeptide is derived from a species belonging to *Arhae*.

40. A method according to claim 33 wherein the second kind of polypeptide is selected from the group consisting of the relB polypeptide derived from *E. coli* K-12 and a functionally equivalent polypeptide.

10

41. A method according to claim 40 wherein the functionally equivalent second kind of polypeptide is derived from the group consisting of a Gram-positive bacterium, a Gram-negative bacterium and *Archae* spp.

15 42. A method according to claim 33 wherein the plasmid contains a gene coding for a gene product of interest.

43. A method according to claim 42 wherein the gene product of interest is an immunologically active gene product.

20

44. A method according to claim 42 wherein the gene product of interest is one that is effective in degradation of an environmental pollutant.

45. A method according to claim 42 wherein the gene product of interest is a pesticidally active product.  
25

46. A method according to claim 45 wherein the gene coding for the pesticidally active gene product is derived from *Bacillus thuringiensis*.

30 47. A method according to claim 33 wherein the microbial cells belong to a Gram-negative bacterial species.

48. A method according to claim 47 wherein the bacterial species is selected from the group consisting of *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp. and *Pseudomonadaceae* spp.

5 49. A method according to claim 33 wherein the microbial cells belong to a Gram-positive bacterial species.

50. A method according to claim 33 wherein the plasmid is one that occurs in the microbial cells at a copy number which is in the range of 1-30 including the range of 1-  
10 10 such as the range of 1-5.

51. A recombinant cell comprising a gene coding for a first kind of polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide and a gene coding for a functionally equivalent polypeptide or a variant or derivative hereof, said  
15 first kind of polypeptide having a toxic effect on the cell, subject to the limitation that when the cell is *E. coli*, the gene coding for the first kind of polypeptide is not derived from *E. coli*.

52. A cell according to claim 51 wherein the cells belong to a bacterial species selected from the group consisting of *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp. and *Pseudomonadaceae* spp.  
20

53. A cell according to claim 51 wherein the cells belong to a Gram-positive bacterial species.

25

54. A cell according to claim 51 wherein the gene coding for *E. coli* K-12 RelE polypeptide or the functional equivalent hereof is located on the chromosome.

55. A cell according to claim 51 wherein the gene coding for the *E. coli* K-12 RelE  
30 polypeptide or the functional equivalent hereof is located on an extrachromosomal replicon.

56. A cell according to claim 51 which comprises at least one gene coding for a gene product of interest.

57. A cell according to claim 56 wherein the gene product of interest is an immunologically active gene product.

58. A cell according to claim 56 wherein the gene product of interest is one that is  
5 effective in degradation of an environmental pollutant.

59. A cell according to claim 51 wherein the gene product of interest is a pesticidally active product.

10 60. A cell according to claim 51 which comprises a gene coding for a second kind of polypeptide that is capable of binding to the first kind of polypeptide whereby the toxic effect hereof is at least reduced.

61. A cell according to claim 51 comprising, operably linked to the gene coding for a  
15 first kind of polypeptide, a regulatable regulatory DNA sequence.

62. A cell according to claim 61 wherein the regulatory DNA sequence regulates the expression of the gene coding for the first kind of polypeptide at the transcriptional level by means of a promoter, the function of which is regulated by the presence or  
20 absence of a chemical compound in the cultivation medium.

63. A cell according to claim 62 wherein the promoter is inducible by a chemical compound.

25 64. A cell according to claim 62 wherein the promoter is suppressible by a first kind of chemical compound and inducible by a second kind of chemical compound whereby, when the first kind of compound is depleted from the medium, the promoter is induced by the second kind of compound.

30 65. A cell according to claim 51 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-negative bacterium.

66. A cell according to claim 65 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-negative bacterium selected from the group



consisting of *Enterobacteriaceae* spp., *Hemophilus* spp., *Vibrionaceae* spp., *Pseudomonadaceae* spp., *Heliobacter* spp. and *Synechosystis* spp.

67. A cell according to claim 51 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-positive bacterium.

68. A cell according to claim 67 wherein the gene coding for the functionally equivalent polypeptide is derived from a Gram-positive bacterium selected from the group consisting of lactic acid bacterial spp., *Bacillaceae* spp. and *Mycobacterium* spp.

10

69. A cell according to claim 68 wherein the Gram-positive bacterium is *Bacillus thuringiensis*.

70. A cell according to claim 51 wherein the gene coding for the functionally equivalent polypeptide is derived from a species belonging to *Arhae*.

71. A cell according to claim 51 which does not contain a gene coding for a second type of polypeptide that is capable of counteracting the cell toxic effect of the first kind of polypeptide.

20

72. A cell according to claim 51 which comprises a gene coding for a second kind of polypeptide that is capable of binding to the first kind of polypeptide, said binding resulting in that the toxic effect of the first kind of polypeptide is at least partially counteracted.

25

73. A cell according to claim 72 wherein the gene coding for the second kind of polypeptide is operably linked to a regulatable regulatory DNA sequence, permitting that the gene coding for said second kind of polypeptide is suppressed under conditions where the gene coding for the RelE polypeptide or the functional equivalent is expressed.

30

74. A cell according to claim 72 wherein the second kind of polypeptide is the RelB polypeptide derived from *E. coli* K-12 or a functionally equivalent polypeptide.

75. A cell according to claim 74 wherein the functionally equivalent second kind of polypeptide is derived from the group consisting of a Gram-positive bacterium, a Gram-negative bacterium and *Archae* spp.

5 76. A cell according to claim 61 wherein the expression of the gene coding for the first kind of polypeptide is stochastically regulated.

77. A cell according to claim 76 wherein the stochastic regulation is effected by operably linking the gene coding for the first kind of polypeptide to a regulatory sequence  
10 that comprises an invertible promoter.

78. A cell according to claim 76 wherein the stochastic regulation is effected by flanking at least part of the regulatory sequence by repeat sequences whereby the at least part of the regulatory sequence is recombinationally excised.

15

79. A composition comprising a cell according to any of claims 51-78.

80. A method of limiting the survival of a cell population in a first or a second environment, which method comprises

20

(i) transforming the cells of said population with a gene coding for a cytotoxic polypeptide, the gene is selected from the group consisting of the gene coding for the *E. coli* K-12 RelE polypeptide, the gene coding for the plasmid F CcdB polypeptide, the gene coding for the plasmid R1 PemK polypeptide, the gene coding for plasmid RP4  
25 ParE polypeptide, the gene coding for the prophage P1 Doc polypeptide and a gene coding for a functionally equivalent polypeptide for anyone of said polypeptides, said gene is expressible in the cells of the population, and operably linked to the gene, a regulatory DNA sequence being regulatable by an environmental factor and which regulates the expression of said gene, and

30

(ii) cultivating the cell population under environmental conditions where the gene coding for the cytotoxic polypeptide is expressed, the expression leading to an at least partial killing of the cell population.

81. A method according to claim 80 wherein the survival of the cell population is limited in a first environment in which the gene is expressed, said cell population thereby being contained in said first environment.

5 82. A method according to claim 80 wherein the survival of the cell population is not limited when present in a first environment, which first environment could change to a second environment physically and/or chemically distinct from the first environment, in which first environment the gene whose expression results in the formation of a cytotoxically active polypeptide is not expressed, but the survival of which cell population  
10 is limited when transferred to a second environment or when present in a physically and/or chemically changed first environment, where the gene is expressed.

83. A method according to claim 80 wherein the survival of a cell population is being limited by providing in the cells a gene coding for a cytotoxic polypeptide which is operably linked to a nucleotide sequence encoding an antitoxin repressor substance  
15 which can undergo a decay when said cells are released to the outer environment to an extent whereby the repressor substance is converted to a non-functional form, whereby as a result of said decay, the function of the cells of the population will be gradually limited.

20

84. A method of containing an extrachromosomal recombinant replicon to a first kind of cell, where said replicon is naturally transferable to a second kind of cell, which method comprises providing on the recombinant extrachromosomal replicon a gene whose expression results in the formation of a cytotoxic polypeptide selected from the  
25 group consisting of the *E. coli* K-12 RelE polypeptide, the plasmid F CcdB polypeptide, the plasmid R1 PemK polypeptide, the plasmid RP4 ParE polypeptide, the prophage P1 Doc polypeptide and a functionally equivalent polypeptide for anyone of said polypeptides to an extent whereby the function of the cell is being limited, said first kind of cells having or being modified to have a chromosomal replicon comprising a regulatory  
30 nucleotide sequence the gene product of which inhibits the expression of said gene or the cell function-limiting effect of the polypeptide and thereby protects said first kind of cells, said regulatory gene being lacking in said second kind of cell, whereby, if a cell of the second kind receives said extrachromosomal recombinant replicon said gene is expressed and has a function-limiting effect on said second kind of cell.

85. A method according to claim 84 wherein the gene product which inhibits the expression of the expression of the gene coding for the polypeptide or the cell function-limiting effect of the polypeptide is selected from the *E. coli* relB polypeptide, the plasmid F CcdA polypeptide, the plasmid R1 PemI polypeptide, the plasmid RP4 ParD  
5 polypeptide, the prophage P1 Phd polypeptide and a functionally equivalent polypeptide of anyone of such polypeptides.

86. A method of stochastically limiting in an environment the survival of a cell population, the method comprising transforming the cells thereof with a recombinant replicon  
10 containing a regulatably expressible gene which, when expressed in a cell encodes a cytotoxic polypeptide selected from the group consisting of the *E. coli* K-12 RelE polypeptide, the plasmid F CcdB polypeptide, the plasmid R1 PemK polypeptide, the plasmid RP4 ParE polypeptide, the prophage P1 Doc polypeptide and a functionally  
equivalent polypeptide for anyone of said polypeptides, the expression of said gene  
15 leading to formation of the polypeptide to an extent whereby the function of the cells is being limited, the expression of said gene is stochastically induced as a result of recombinational excision of an excisable negatively functioning regulatory nucleotide sequence which, while present in the cells, inhibits expression of the gene coding for the  
polypeptide, said negatively functioning regulatory nucleotide sequence being con-  
20 tained in the recombinant replicon or in an other recombinant replicon present in cells of the population containing the replicon.

87. A cell according to claim 51 that is an animal cell.

25 88. A cell according to claim 87 that is selected from the group consisting of a mammal cell, a human cell and an insect cell.

89. A method according to any one of claims 1, 26, 32, 80, 84 or 86 wherein the first kind of polypeptide inhibits translation.

1/22



Plasmids	relBE derived fragments	$\beta$ -galactosidase activity	Fold of repression
pKG4001		1250	1
pKG4002		28	45

Fig. 1

2/22



Fig. 2

3/22




Fig. 3

4/22




Fig. 4



5/22




Fig. 5





Fig. 6A





Fig. 6B

8/22




Fig. 7

9/22




Fig. 8




Fig. 9

11/22

DNA sequence of the *relB*<sub>sp2</sub> locus of *S. pneumoniae* (reverse complement of part of contig 80 in the tigr database)

```

  1  ACCCATGAGC ATTCAGATCA TATCCATGGA GTAGGCGTTT TGGCTCGCAA
 51  GTATGGTATG GATCTTTATG CCAATGAAAA GACCTGGCAA GCTATGAAAA
101  ATAGTAAATA TCTTGGCAAG GTGGATTCTT CGCAAAAGCA TATTTTTGAA
151  ATGGGTAAAA CCAAACCTT TGGAGATATC GACATCGAGA GTTTTGGTGT
201  AAGCCATGAT GCAGTCGCAC CGCAGTTCTA TCGCTTTATG AAGGATGATA
251  AGAGTTTTGT CCTCTTGACA GATACAGGTT ATGTCAGTGA CCGTATGGCG
301  GGCATTGTCTG AAAATGCGGA TGGTTATCTT ATCGAGGCCA ACCATGATGT
351  AGAGATTTTG CGATCAGGTT CTTACGCTTG GCGACTCAAA CAACGAATCC
401  TATCTGACCT TGGTCACCTT TCTAACGAGG ACGGTGCTGA AGCTATGATT
451  CGGACGCTAG GAAATCGTAC TAAGAAGATT TACCTTGGGC ATTTATCTAA
501  GGAAAACAAT ATCAAGGAAC TGGCTCATAT GACCATGGTC AATCAGCTGG
551  CTCAAGCTGA TCTGGGAGTC GGAGTAGACT TTAAGGTTTA TGATACCTCA
601  CCAGATACCG CAACACCATT GACAGAGATA TAGAAAGAAC GCTGAGAAGG
651  TGTTCTTTTT ATATTGACTG AACACCTAAA AAGTAATACA ATGGTGTTAC
      SD                               Start relBsp2
701  CATTAAAAAA GGGAGCACAA AAGATGACTA CTATTACATT AAAAGTTTCT
751  GAAGCTGATA AAACATTTAT GAAAGCAATG GCTAAGTTTG AAGGAGTTTC
801  CCTTTCGGAA CTTATTTCGA CCAAACTCT TGAAGCCCTA GAAGATGAAT
851  ACGATGCTCG TGTGGCAGAT TTAGCCTATC AAGAGTATTT AGAAGACTTG
      SD
901  GAAAAAGGAG TTGAACCCAT TACTTGGGAA GAAATGATGC ATGATTTAGG
      Start relEsp2/End relBsp2
951  CTTGAAGGAT GAATAATTTG TATAAATTAG TTCCAACAAG ACGTTTTATC
1001 AAGCAATTGA AAAAATTGGA CCGTTATACG CAGAAGCTAA TTACAAACTA
1051 TTTACAAACC AATGTTTTGG AAGACCCAAG ACGACACGGA AAGGCTTTGG
1101 TTGGTAATCG CGTTGGTCAA TGGCGCTATA GAATTGGTAA TTATCGAGTT
1151 ATCGTACAAA TTGTAGATGA TGAATTAGTC GTTGCTACTC TAGAAGTTGG
      End relEsp2
1201 TCATCGGAGA GATATTTATT GAATTACTTT TTTCAAGTAA ATTTTCATTC
1251 TTTAAAATAT GATATAATAG GGCTTACAAT ATAAGAGAGG TTGCTGATGA
1301 CAATAGATAT TAGTGAAGAA AGTTTGGCTA AAGAATCGGC TGAATTATTA
1351 AAAATTCCTT TAAAAGATCG AACGACAAAG AAATCAATTG TTTGGGCTAC
1401 GCATTCTTAT GAATTGTTAG GAAAGGGATT TGCTCCAAGT GATCGCATCA
1451 ATCCAAGTAA GGTTACTGGA AACTTTGCCA ACTTGATTCA GCCCAGATCT
1501 GAAAAATCCA AATATGAACA AAAAGATCGG ACAAAAATAA GGGCAGAAGT

```

#### Protein sequences:

*RelB*<sub>sp2</sub> (pHi=4.2):

mttitlkvse adktfmkama kfegvslsel irtktleale deydarvadl aygeyledle  
kgvepitwee mmhdlglkde

*RelE*<sub>sp2</sub> (pHi=11.1):

mnnyklvpt rrfikqlkkl drytqklitn yltqnvledp rrrhgkalvgn  
rvqqwryrig nyrvivqivd delvvatlev ghrrdiy

Fig. 10

12/22




Fig. 11



13/22




Fig. 12

14/22




Fig. 13

15/22




Fig. 14

16/22




Fig. 15

17/22




Fig. 16

18/22




Fig. 17

19/22




Fig. 18




Fig. 19



21/22




Fig. 20

22/22




Fig. 21

## SEQUENCE LISTING

5       <110> GX BioSystems A/S  
          Gerdes, Kenn  
          Gotfredsen, Marie  
          Grønland, Hugo  
          Pedersen, Kim  
10       Kristoffersen, Peter  
  
          <120> Cytotoxin-based biological containment  
  
15       <130> 20935 PC 1  
  
          <150> PA 1998 00627  
          <151> 1998-05-07  
  
20       <150> US 60/085,067  
          <151> 1998-05-12  
  
          <160> 59  
  
25       <170> FastSEQ for Windows Version 3.0  
  
          <210> 1  
          <211> 51  
          <212> DNA  
30       <213> Artificial Sequence  
  
          <220>  
          <223> Primer relE1B was used for the amplification of  
                  relEK-12 by PCR on pBD2430  
35       <400> 1  
          ccccg gatcc ataaggagtt ttataaatgg cgtattttct ggattttgac g 51  
  
          <210> 2  
40       <211> 38  
          <212> DNA  
          <213> Artificial Sequence  
  
          <220>  
45       <223> Primer relE1B was used for the amplification of  
                  relEK-12 by PCR on pBD2430  
  
          <400> 2  
          ccccctcga ggtcgactca gagaatgcgt ttgaccgc 38  
50  
  
          <210> 3  
          <211> 28  
          <212> DNA  
          <213> Artificial Sequence  
55  
  
          <220>  
          <223> Primer RelB-P307/1 was used for the generation of  
                  a PCR-fragment from pNZ945

<400> 3  
cccccgatc ccagtcttga aaggtggc 28

5 <210> 4  
<211> 29  
<212> DNA  
<213> Artificial Sequence

10 <220>  
<223> Primer RelB-P307/2 was used for the generation of  
a PCR-fragment from pNZ945

<400> 4  
15 cccccgaatt ctcataggta tttatccag 29

<210> 5  
<211> 27  
<212> DNA  
<213> Artificial Sequence

20 <220>  
<223> Primer relE-p307/3 was used to PCR-amplify the  
gene relEP307 from pNZ945

25 <400> 5  
cccccgatcc agatctggat aaatacc 27

<210> 6  
<211> 32  
30 <212> DNA  
<213> Artificial Sequence

<220>  
35 <223> Primer relE-P307/2 was used to PCR-amplify the  
gene relEP307 from pNZ945

<400> 6  
cccccggaatt cgtaactttc tgtgtttatt gc 32

40 <210> 7  
<211> 28  
<212> DNA  
<213> Artificial Sequence

45 <220>  
<223> Primer relE-P307/4 was used for the generation of  
a DNA fragment encoding relEP307 by PCR

<400> 7  
50 cccccgagct cagatctgga taaatacc 28

<210> 8  
<211> 32  
55 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer relE-P307/5 was used for the generation of  
a DNA fragment encoding relEP307 by PCR

	<400> 8		
	ccccgcgatg cgtaactttc tgtgtttatt gc		32
5	<210> 9		
	<211> 1461		
	<212> DNA		
	<213> E. coli K-12		
10	<400> 9		
	cttaatttca ggccccatcg gatcacacat ggagagtttt tatgaataac cccgtctgtc 60		
	ttgatgactg gttgattggc tttaaaagct tgttgacagg ggtaaacggt cggcaataat 120		
	tttctgccgc atgcggggtgt tgcataaaac gtgttacggt cctttatcga caggtcagggt 180		
	caccgctcac ccgccgacga gaaagcaaca ctgacatgct aaagcaaaaa atagatgaat 240		
15	aagttgagtt gtgcatatgt agcctgaccg tcacaaagta tatgggtgtct gtaccagtaa 300		
	gatgatggcc ggactcttta aaaacgagct gacctgcaca atacaggatg gacttagcaa 360		
	tggctgctcc tggcacaaag cggacagtga tcaccgttct tacgactact ttctgacttc 420		
	cttcgtgact tgccttaagc atgttgtagt rbmrnarbst artgcgatac ttgtaatgac 480		
	atgtgtaatt acaagagggtg taagacatgg gtargcatta acctgcgtat tgacgatgaa 540		
20	cttaaagcgc gttcttacgc cgcgcttgaa aaaatgggtg taactccttc tgaagcgctt 600		
	cgtctcatgc tcgagtatat cgctgacaat gaacgcttgc cgttcaaaca gacactcctg 660		
	agtgatgaag atgctgaact tgtggagata gtgaaagaac ggcttcgtaa tcctndrbst 720		
	artraagcca gtacgtgtga cgctggatga actctgatgg cgtattttct ggattttgac 780		
	gagcgggcac taaaggaatg gcgaaagctg ggctcgacgg tacgtgaaca gttgaaaaag 840		
25	aagctggttg aagtacttga gtcaccccg attgaagcaa acaagctccg tggtagcct 900		
	gattgttaca agattaagct ccggtcttca ggctatcgcc ttgtatacca ggttatagac 960		
	gagaaagttg tcgttttctg gattttctgtt gggaaaagag aacgctcgga agtatatagc 1020		
	gaggndrcgg tcaaacgc atctctgaacc aaagcatgac atctctgttt cgcaccgsta 1080		
	rthkcragg tgacacttct gctttgcgtt gacaggagaa gcaggctatg aagcagcaaa 1140		
30	aggcgatggt aatcgccctg atcgctcatct gtttaaccgt catagtgaag gcactggtaa 1200		
	cgaggaaaga cctctgcgag gtacgaatcc gaaccgndhk caccagacgg aggtcgctgt 1260		
	cttcacagct tacgaacctg aggagtaaga gacccggcgg gggagaaatc cctcgccacc 1320		
	tctgatgtgg caggcatcct caacgcaccc gcacttaacc cgcttcggcg ggtttttggt 1380		
	tttattttca arttcgcgtt tgaagttctg gacggtgccg gaatagaatc aaaaatactt 1440		
35	aagtdataba saccsnnumb r		1461
	<210> 10		
	<211> 88		
	<212> PRT		
40	<213> Bacterium		
	<220>		
	<223> protein RelE-Mj2		
45	<400> 10		
	Met Lys Val Leu Phe Ala Lys Thr Phe Val Lys Asp Leu Lys His Val		
	1 5 10 15		
	Pro Gly His Ile Arg Lys Arg Ile Lys Leu Ile Ile Glu Glu Cys Gln		
	20 25 30		
50	Asn Ser Asn Ser Leu Asn Asp Leu Lys Leu Asp Ile Lys Lys Ile Lys		
	35 40 45		
	Gly Tyr His Asn Tyr Tyr Arg Ile Arg Val Gly Asn Tyr Arg Ile Gly		
	50 55 60		
	Ile Glu Val Asn Gly Asp Thr Ile Ile Phe Arg Arg Val Leu His Arg		
55	65 70 75 80		
	Lys Ser Ile Tyr Asp Tyr Phe Pro		
	85		

<210> 11  
 <211> 91  
 <212> PRT  
 <213> Bacterium  
 5  
 <220>  
 <223> protein RelE-Mj3  
 <400> 11  
 10 Met Lys Gln Trp Lys Tyr Leu Leu Lys Lys Ser Phe Ile Lys Asp Leu  
 1 5 10 15  
 Lys Glu Leu Pro Lys Asn Ile Gln Glu Lys Ile Lys Lys Leu Val Phe  
 20 25 30  
 Glu Glu Ile Pro Asn Lys Asn Asn Pro Pro Glu Ile Pro Asn Val Lys  
 15 35 40 45  
 Lys Leu Lys Gly Ala Asp Ser Tyr Tyr Arg Ile Arg Val Gly Asp Tyr  
 50 55 60  
 Arg Ile Gly Phe Lys Tyr Glu Asn Gly Lys Ile Val Phe Tyr Arg Val  
 65 70 75 80  
 20 Leu His Arg Lys Gln Ile Tyr Lys Arg Phe Pro  
 85 90  
 <210> 12  
 <211> 87  
 25 <212> PRT  
 <213> Bacterium  
 <220>  
 <223> protein RelE-Af1  
 30  
 <400> 12  
 Met Phe Arg Val Val His Arg Lys Ala Thr Gln Glu Leu Lys Arg  
 1 5 10 15  
 Leu Lys Lys Ala His Leu Lys Lys Phe Gly Val Leu Leu Glu Thr Leu  
 20 25 30  
 Lys Thr Asp Pro Ile Pro Trp Lys Arg Phe Asp Val Lys Lys Ile Glu  
 35 40 45  
 Gly Glu Glu Asn Thr Tyr Arg Ile Arg Ile Gly Asp Phe Arg Val Ile  
 50 55 60  
 40 Tyr Phe Leu Asp Lys Pro Thr Lys Thr Val His Ile Leu Lys Val Glu  
 65 70 75 80  
 Arg Arg Gly Lys Val Tyr Asp  
 85  
 45 <210> 13  
 <211> 90  
 <212> PRT  
 <213> Bacterium  
 50  
 <220>  
 <223> protein RelE-Mj1  
 <400> 13  
 55 Met Lys Phe Asn Val Glu Ile His Lys Arg Val Leu Lys Asp Leu Lys  
 1 5 10 15  
 Asp Leu Pro Pro Ser Asn Leu Lys Lys Phe Lys Glu Leu Ile Glu Thr  
 20 25 30  
 Leu Lys Thr Asn Pro Ile Pro Lys Glu Lys Phe Asp Ile Lys Arg Leu  
 35 40 45

Lys Gly Ser Asp Glu Val Tyr Arg Val Arg Ile Gly Lys Phe Arg Val  
 50 55 60  
 Gln Tyr Val Val Leu Trp Asp Asp Arg Ile Ile Ile Arg Lys Ile  
 65 70 75 80  
 5 Ser Arg Arg Glu Gly Ala Tyr Lys Asn Pro  
 85 90

<210> 14  
 <211> 74  
 10 <212> PRT  
 <213> Bacterium

<220>  
 <223> protein RelE-Bt  
 15

<400> 14  
 Met Lys Phe Ile Ala Lys Gln Glu Lys Gly Ile Gln Lys Arg Ile Ala  
 1 5 10 15  
 Glu Gly Leu Lys Gly Leu Leu Lys Ile Pro Pro Glu Gly Asp Ile Lys  
 20 20 25 30  
 Ser Met Lys Gly Tyr Thr Glu Leu Tyr Arg Leu Arg Ile Gly Thr Phe  
 35 40 45  
 Arg Ile Leu Phe Glu Ile Asn His Asp Glu Lys Val Ile Tyr Ile Gln  
 50 55 60  
 25 Ala Ile Gly Asn Arg Gly Asp Ile Tyr Lys  
 65 70

<210> 15  
 <211> 95  
 30 <212> PRT  
 <213> Bacterium

<220>  
 <223> protein RelE-P307  
 35

<400> 15  
 Met Arg Tyr Gln Val Lys Phe Arg Glu Asp Ala Leu Lys Glu Trp Gln  
 1 5 10 15  
 Lys Leu Asp Lys Ala Ile Gln Gln Gln Phe Ala Lys Lys Leu Lys Lys  
 40 20 25 30  
 Cys Cys Asp Asn Pro His Ile Pro Ser Ala Lys Leu Arg Gly Ile Lys  
 35 40 45  
 Asp Cys Tyr Lys Ile Lys Leu Arg Ala Ser Gly Phe Arg Leu Val Tyr  
 50 55 60  
 45 Gln Val Ile Asp Glu Gln Leu Ile Ile Ala Val Val Ala Val Gly Lys  
 65 70 75 80  
 Arg Glu Arg Ser Asp Val Tyr Asn Leu Ala Ser Glu Arg Met Arg  
 85 90 95

50 <210> 16  
 <211> 95  
 <212> PRT  
 <213> Bacterium

55 <220>  
 <223> protein RelEK-12

<400> 16  
 Met Ala Tyr Phe Leu Asp Phe Asp Glu Arg Ala Leu Lys Glu Trp Arg  
 1 5 10 15  
 Lys Leu Gly Ser Thr Val Arg Glu Gln Leu Lys Lys Lys Leu Val Glu  
 20 25 30  
 Val Leu Glu Ser Pro Arg Ile Glu Ala Asn Lys Leu Arg Gly Met Pro  
 35 40 45  
 Asp Cys Tyr Lys Ile Lys Leu Arg Ser Ser Gly Tyr Arg Leu Val Tyr  
 50 55 60  
 Gln Val Ile Asp Glu Lys Val Val Val Phe Val Ile Ser Val Gly Lys  
 65 70 75 80  
 Arg Glu Arg Ser Glu Val Tyr Ser Glu Ala Val Lys Arg Ile Leu  
 85 90 95

15 <210> 17  
 <211> 96  
 <212> PRT  
 <213> Bacterium

20 <220>  
 <223> protein RelE-Vc

<400> 17  
 Met Thr Tyr Lys Leu Glu Phe Lys Lys Ser Ala Leu Lys Glu Trp Lys  
 1 5 10 15  
 Lys Leu Ala Val Pro Leu Gln Gln Gln Phe Lys Lys Lys Leu Ile Glu  
 20 25 30  
 Arg Leu Glu Asn Pro His Val Pro Ser Ala Lys Leu Ser Gly Ala Glu  
 35 40 45  
 Asn Ile Tyr Lys Ile Lys Leu Arg Gln Ser Gly Tyr Arg Leu Val Tyr  
 50 55 60  
 Gln Val Glu Asn Asp Ile Val Val Thr Val Leu Ala Val Gly Lys  
 65 70 75 80  
 Arg Glu Arg Ser Glu Val Tyr Thr Lys Ala Leu Gln Arg Leu Asp Asp  
 85 90 95

40 <210> 18  
 <211> 87  
 <212> PRT  
 <213> Bacterium

<220>  
 <223> protein RelE-Mtl

45 <400> 18  
 Met Pro Tyr Thr Val Arg Phe Thr Thr Thr Ala Arg Arg Asp Leu His  
 1 5 10 15  
 Lys Leu Pro Pro Arg Ile Leu Ala Ala Val Val Glu Phe Ala Phe Gly  
 20 25 30  
 Asp Leu Ser Arg Glu Pro Leu Arg Val Gly Lys Pro Leu Arg Arg Glu  
 35 40 45  
 Leu Ala Gly Thr Phe Ser Ala Arg Arg Gly Thr Tyr Arg Leu Leu Tyr  
 50 55 60  
 Arg Ile Asp Asp Glu His Thr Thr Val Val Ile Leu Arg Val Asp His  
 55 65 70 75 80  
 Arg Ala Asp Ile Tyr Arg Arg  
 85



<210> 19  
 <211> 97  
 <212> PRT  
 <213> Bacterium

5

<220>  
 <223> protein RelE-Mt2

<400> 19  
 10 Met Ser Asp Asp His Pro Tyr His Val Ala Ile Thr Ala Thr Ala Ala  
     1                    5                    10                    15  
     Arg Asp Leu Gln Arg Leu Pro Glu Lys Ile Ala Ala Ala Cys Val Glu  
             20                    25                    30  
 15 Phe Val Phe Gly Pro Leu Leu Asn Asn Pro His Arg Leu Gly Lys Pro  
             35                    40                    45  
     Leu Arg Asn Asp Leu Glu Gly Leu His Ser Ala Arg Arg Gly Asp Tyr  
             50                    55                    60  
     Arg Val Val Tyr Ala Ile Asp Asp Gly His His Arg Val Glu Ile Ile  
             65                    70                    75                    80  
 20 His Ile Ala Arg Arg Ser Ala Ser Tyr Arg Met Asn Pro Cys Arg Pro  
                     85                    90                    95  
     Arg

25

<210> 20  
 <211> 102  
 <212> PRT  
 <213> Bacterium

30

<220>  
 <223> protein RelE-Hi

<400> 20  
 35 Met Ser Glu Glu Lys Pro Leu Lys Val Ser Tyr Ser Lys Gln Phe Val  
     1                    5                    10                    15  
     Arg Asp Leu Thr Asp Leu Ala Lys Arg Ser Pro Asn Val Leu Ile Gly  
             20                    25                    30  
     Ser Lys Tyr Ile Thr Ala Ile His Cys Leu Leu Asn Arg Leu Pro Leu  
             35                    40                    45  
 40 Pro Glu Asn Tyr Gln Asp His Ala Leu Val Gly Glu Trp Lys Gly Tyr  
     50                    55                    60  
     Arg Asp Cys His Ile Gln Gly Asp Leu Val Leu Ile Tyr Gln Tyr Val  
             65                    70                    75                    80  
     Ile Gln Asp Glu Phe Asp Glu Leu Lys Phe Ser Arg Leu Asn Ile His  
             85                    90                    95  
 45 Ser Gln Thr Ala Leu Lys  
             100

50

<210> 21  
 <211> 92  
 <212> PRT  
 <213> Bacterium

55

<220>  
 <223> protein RelEk12sos

<400> 21  
 Met Ile Gln Arg Asp Ile Glu Tyr Ser Gly Gln Tyr Ser Lys Asp Val  
 1 5 10 15  
 Lys Leu Ala Gln Lys Arg His Lys Asp Met Asn Lys Leu Lys Tyr Leu  
 5 20 25 30  
 Met Thr Leu Leu Ile Asn Asn Thr Leu Pro Leu Pro Ala Val Tyr Lys  
 35 40 45  
 Asp His Pro Leu Gln Gly Ser Trp Lys Gly Tyr Arg Asp Ala His Val  
 50 55 60  
 10 Glu Pro Asp Trp Ile Leu Ile Tyr Lys Leu Thr Asp Lys Leu Leu Arg  
 65 70 75 80  
 Phe Glu Arg Thr Gly Thr His Ala Ala Leu Phe Gly  
 85 90

15 <210> 22  
 <211> 88  
 <212> PRT  
 <213> Bacterium

20 <220>  
 <223> protein RelE-Hp

<400> 22  
 Met Leu Lys Leu Asn Leu Lys Lys Ser Phe Gln Lys Asp Phe Asp Lys  
 1 5 10 15  
 Leu Leu Leu Asn Gly Phe Asp Asp Ser Val Leu Asn Glu Val Ile Leu  
 20 25 30  
 Thr Leu Arg Lys Lys Glu Pro Leu Asp Pro Gln Phe Gln Asp His Ala  
 35 40 45  
 30 Leu Lys Gly Lys Trp Lys Pro Tyr Arg Glu Cys His Ile Lys Pro Asp  
 50 55 60  
 Val Leu Leu Val Tyr Leu Val Lys Asp Asp Glu Leu Ile Leu Leu Arg  
 65 70 75 80  
 Leu Gly Ser His Ser Glu Leu Phe  
 85

35 <210> 23  
 <211> 92  
 <212> PRT  
 <213> Bacterium

40 <220>  
 <223> protein RelE-Af2

45 <400> 23  
 Met Ala Trp Lys Val Arg Tyr His Lys Lys Ala Ile Lys Phe Leu Glu  
 1 5 10 15  
 Lys Leu Asp Glu Gly Lys Arg Ser Ile Leu Leu Ser Lys Ile Gln Glu  
 20 25 30  
 50 Leu Val Asn Ser Leu Glu Ser Gly Val Leu Pro Ile Gln Arg Met Asp  
 35 40 45  
 Ile Lys Arg Leu Lys Gly Val Trp Asp Gly Phe Leu Arg Leu Arg Val  
 50 55 60  
 Gly Glu Val Arg Ile Ile Phe Lys Ile Asn Val Glu Asp Glu Thr Ile  
 55 65 70 75 80  
 Phe Ile Tyr Ser Ile His Phe Arg Glu Lys Val Tyr  
 85 90

5  
 <210> 24  
 <211> 86  
 <212> PRT  
 <213> Bacterium

<220>  
 <223> protein RelE-Af4  
 <400> 24  
 10 Met Asn Glu Val Leu Ile His Lys Lys Phe Leu Asp Gly Leu Asp Ser  
 1 5 10 15  
 Gly Arg Arg Ser Lys Val Leu Asp Ala Ile Arg Met Leu Lys Asp Phe  
 20 25 30  
 Pro Ile Ile Arg Ala Asp Ile Lys Lys Ile Gly Pro Lys Thr Tyr Arg  
 15 35 40 45  
 Leu Arg Lys Gly Glu Ile Arg Ile Ile Phe Asp Phe Asp Ile Gly Thr  
 50 55 60  
 Asn Arg Val Phe Val Lys Phe Ala Ala Ser Glu Gly Val Phe Thr Lys  
 65 70 75 80  
 20 Thr Glu Glu Lys Phe Phe  
 85

25  
 <210> 25  
 <211> 85  
 <212> PRT  
 <213> Bacterium

<220>  
 <223> protein RelE-Af3  
 30  
 <400> 25  
 Met Asn Tyr Lys Ala Gln Phe Ser Glu Glu Phe Leu Lys Ile Ala Lys  
 1 5 10 15  
 Lys Leu Lys Glu Lys Asp Pro Glu Leu Leu Lys Arg Leu Gln Ser Lys  
 35 20 25 30  
 Val Glu Glu Ile Ile Lys Gln Pro Glu His Tyr Lys Pro Leu Arg Gly  
 35 40 45  
 Gln Met Lys Gly Leu Arg Arg Ala His Val Gly Lys Phe Val Ile Ile  
 50 55 60  
 40 Phe Lys Val Glu Glu Asp Thr Val Lys Phe Val Thr Phe Lys His His  
 65 70 75 80  
 Asn His Ala Tyr Lys  
 85

45  
 <210> 26  
 <211> 120  
 <212> PRT  
 <213> Bacterium

50  
 <220>  
 <223> protein RelE-Sy

<400> 26  
 55 Met Ser Asn Asn Leu His Leu Val Asn Ile Asp Phe Thr Pro Glu Tyr  
 1 5 10 15  
 Arg Arg Ser Leu Lys Tyr Leu Ala Lys Lys Tyr Arg Asn Ile Arg Ser  
 20 25 30  
 Asp Val Gln Pro Ile Ile Glu Ala Leu Gln Lys Gly Val Ile Ser Gly  
 35 40 45

10

Asp Arg Leu Ala Gly Phe Gly Ser Asp Ile Tyr Val Tyr Lys Leu Arg  
 50 55 60  
 Ile Lys Asn Ser Asn Ile Gln Lys Gly Lys Ser Ser Gly Tyr Arg Leu  
 65 70 75 80  
 5 Ile Tyr Leu Leu Glu Ser Glu Asn Ser Ile Leu Leu Leu Thr Ile Tyr  
 85 90 95  
 Ser Lys Ala Glu Gln Glu Asp Ile Ala Ala Ser Asp Ile Asn Ser Ile  
 100 105 110  
 10 Leu Gly Glu Tyr Ser Ile Glu Asp  
 115 120

&lt;210&gt; 27

&lt;211&gt; 86

&lt;212&gt; PRT

15 &lt;213&gt; Bacterium

&lt;220&gt;

&lt;223&gt; protein RelB-SOS

20 &lt;400&gt; 27

Met Ala Ala Asn Ala Phe Val Arg Ala Arg Ile Asp Glu Asp Leu Lys  
 1 5 10 15  
 Asn Gln Ala Ala Asp Val Leu Ala Gly Met Gly Leu Thr Ile Ser Asp  
 20 25 30  
 25 Leu Val Arg Ile Thr Leu Thr Lys Val Ala Arg Glu Lys Ala Leu Pro  
 35 40 45  
 Phe Asp Leu Arg Glu Pro Asn Gln Leu Thr Ile Gln Ser Ile Lys Asn  
 50 55 60  
 Ser Glu Ala Gly Ile Asp Val His Lys Ala Lys Asp Ala Asp Asp Leu  
 65 70 75 80  
 30 Phe Asp Lys Leu Gly Ile  
 85

&lt;210&gt; 28

35 &lt;211&gt; 82

&lt;212&gt; PRT

&lt;213&gt; Bacterium

&lt;220&gt;

40 &lt;223&gt; protein RelB-VC

&lt;400&gt; 28

Met Thr Thr Arg Ile Leu Ala Asp Val Ala Ala Ser Ile Thr Glu Phe  
 1 5 10 15  
 45 Lys Ala Asn Pro Met Lys Val Ala Thr Ser Ala Phe Gly Ala Pro Val  
 20 25 30  
 Ala Val Leu Asn Arg Asn Glu Pro Ala Phe Tyr Cys Val Pro Ala Ser  
 35 40 45  
 Thr Tyr Glu Ile Met Met Asp Lys Leu Glu Asp Leu Glu Leu Leu Ala  
 50 55 60  
 50 Ile Ala Lys Glu Arg Leu Ser Glu Asp Ser Val Ser Val Asn Ile Asp  
 65 70 75 80  
 Asp Leu

55

<210> 29  
 <211> 83  
 <212> PRT  
 <213> Bacterium

5

<220>  
 <223> protein RelB

<400> 29  
 10 Met Pro Asn Ile Ile Leu Ser Asp Thr Ser Ala Ser Val Ser Glu Leu  
 1 5 10 15  
 Lys Lys Asn Pro Met Ala Thr Val Ser Ala Gly Asp Gly Phe Pro Val  
 20 25 30  
 Ala Ile Leu Asn Arg Asn Gln Pro Ala Phe Tyr Cys Val Pro Ala Glu  
 15 35 40 45  
 Leu Tyr Glu Lys Met Leu Asp Ala Leu Asp Asp Gln Glu Leu Val Lys  
 50 55 60  
 Leu Val Ala Glu Arg Ser Asn Gln Pro Leu His Asp Val Asp Leu Asp  
 65 70 75 80  
 20 Lys Tyr Leu

<210> 30  
 <211> 93  
 <212> PRT  
 <213> Bacterium

25

<220>  
 <223> protein RelB-Mt1

30

<400> 30  
 Met Arg Ile Leu Pro Ile Ser Thr Ile Lys Gly Lys Leu Asn Glu Phe  
 1 5 10 15  
 Val Asp Ala Val Ser Ser Thr Gln Asp Gln Ile Thr Ile Thr Lys Asn  
 35 20 25 30  
 Gly Ala Pro Ala Ala Val Leu Val Gly Ala Asp Glu Trp Glu Ser Leu  
 35 40 45  
 Gln Glu Thr Leu Tyr Trp Leu Ala Gln Pro Gly Ile Arg Glu Ser Ile  
 50 55 60  
 40 Ala Glu Ala Asp Ala Asp Ile Ala Ser Gly Arg Thr Tyr Gly Glu Asp  
 65 70 75 80  
 Glu Ile Arg Ala Glu Phe Gly Val Pro Arg Arg Pro His  
 85 90

<210> 31  
 <211> 89  
 <212> PRT  
 <213> Bacterium

45

<220>  
 <223> protein RelB-Mt2

50

<400> 31  
 Met Ala Val Val Pro Leu Gly Glu Val Arg Asn Arg Leu Ser Glu Tyr  
 1 5 10 15  
 Val Ala Glu Val Glu Leu Thr His Glu Arg Ile Thr Ile Thr Arg His  
 20 25 30  
 Gly His Pro Ala Ala Val Leu Ile Ser Ala Asp Asp Leu Ala Ser Ile  
 35 40 45

12

Glu Glu Thr Leu Glu Val Leu Arg Thr Pro Gly Ala Ser Glu Ala Ile  
     50                    55                    60  
 Arg Glu Gly Leu Ala Asp Val Ala Ala Gly Arg Phe Val Ser Asn Asp  
     65                    70                    75                    80  
 5 Glu Ile Arg Asn Arg Tyr Thr Ala Arg  
                     85

<210> 32  
 <211> 97  
 10 <212> PRT  
     <213> Bacterium

<220>  
 <223> protein RelB-K12-2  
 15

<400> 32  
 Met His Arg Ile Leu Ala Glu Lys Ser Val Asn Ile Thr Glu Leu Arg  
     1                    5                    10                    15  
 20 Lys Asn Pro Ala Lys Tyr Phe Ile Asp Gln Pro Val Ala Val Leu Ser  
                     20                    25                    30  
 Asn Asn Arg Pro Ala Gly Tyr Leu Leu Ser Ala Ser Ala Phe Glu Ala  
                     35                    40                    45  
 Leu Met Asp Met Leu Ala Glu Gln Glu Glu Lys Lys Pro Ile Lys Ala  
     50                    55                    60  
 25 Arg Phe Arg Pro Ser Ala Ala Arg Leu Glu Glu Ile Thr Arg Arg Ala  
     65                    70                    75                    80  
 Glu Gln Tyr Leu Asn Asp Met Thr Asp Asp Asp Phe Asn Asp Phe Lys  
                     85                    90                    95  
 Glu

30  
 <210> 33  
 <211> 68  
 <212> PRT  
 35 <213> Bacterium

<220>  
 <223> protein RelB-St

40 <400> 33  
 Met Phe Met Arg Thr Val Asn Tyr Ser Glu Ala Arg Gln Asn Leu Ala  
     1                    5                    10                    15  
 Glu Val Leu Glu Ser Ala Val Thr Gly Gly Pro Val Thr Ile Thr Arg  
                     20                    25                    30  
 45 Arg Gly His Lys Ser Ala Val Ile Ile Ser Ala Glu Glu Phe Glu Arg  
                     35                    40                    45  
 Tyr Gln Thr Ala Arg Met Asp Asp Glu Phe Ala Ala Ile Met Ala Val  
     50                    55                    60  
 His Gly Asn Glu

50 65  
 <210> 34  
 <211> 79  
 <212> PRT  
 55 <213> Bacterium

<220>  
 <223> protein RelB-coli

<400> 34  
 1 Met Gly Ser Ile Asn Leu Arg Ile Asp Asp Glu Leu Lys Ala Arg Ser  
 5 Tyr Ala Ala Leu Glu Lys Met Gly Val Thr Pro Ser Glu Ala Leu Arg  
 10 Leu Met Leu Glu Tyr Ile Ala Asp Asn Glu Arg Leu Pro Phe Lys Gln  
 15 Thr Leu Leu Ser Asp Glu Asp Ala Glu Leu Val Glu Ile Val Lys Glu  
 20 Arg Leu Arg Asn Pro Lys Pro Val Arg Val Thr Leu Asp Glu Leu  
 25  
 30  
 35  
 40  
 45  
 50  
 55  
 60  
 65  
 70  
 75  
 80  
 85  
 90  
 95  
 100  
 105  
 110  
 115  
 120  
 125  
 130  
 135  
 140  
 145  
 150  
 155  
 160  
 165  
 170  
 175  
 180  
 185  
 190  
 195  
 200  
 205  
 210  
 215  
 220  
 225  
 230  
 235  
 240  
 245  
 250  
 255  
 260  
 265  
 270  
 275  
 280  
 285  
 290  
 295  
 300  
 305  
 310  
 315  
 320  
 325  
 330  
 335  
 340  
 345  
 350  
 355  
 360  
 365  
 370  
 375  
 380  
 385  
 390  
 395  
 400  
 405  
 410  
 415  
 420  
 425  
 430  
 435  
 440  
 445  
 450  
 455  
 460  
 465  
 470  
 475  
 480  
 485  
 490  
 495  
 500  
 505  
 510  
 515  
 520  
 525  
 530  
 535  
 540  
 545  
 550  
 555  
 560  
 565  
 570  
 575  
 580  
 585  
 590  
 595  
 600  
 605  
 610  
 615  
 620  
 625  
 630  
 635  
 640  
 645  
 650  
 655  
 660  
 665  
 670  
 675  
 680  
 685  
 690  
 695  
 700  
 705  
 710  
 715  
 720  
 725  
 730  
 735  
 740  
 745  
 750  
 755  
 760  
 765  
 770  
 775  
 780  
 785  
 790  
 795  
 800  
 805  
 810  
 815  
 820  
 825  
 830  
 835  
 840  
 845  
 850  
 855  
 860  
 865  
 870  
 875  
 880  
 885  
 890  
 895  
 900  
 905  
 910  
 915  
 920  
 925  
 930  
 935  
 940  
 945  
 950  
 955  
 960  
 965  
 970  
 975  
 980  
 985  
 990  
 995  
 1000  
 1005  
 1010  
 1015  
 1020  
 1025  
 1030  
 1035  
 1040  
 1045  
 1050  
 1055  
 1060  
 1065  
 1070  
 1075  
 1080  
 1085  
 1090  
 1095  
 1100  
 1105  
 1110  
 1115  
 1120  
 1125  
 1130  
 1135  
 1140  
 1145  
 1150  
 1155  
 1160  
 1165  
 1170  
 1175  
 1180  
 1185  
 1190  
 1195  
 1200  
 1205  
 1210  
 1215  
 1220  
 1225  
 1230  
 1235  
 1240  
 1245  
 1250  
 1255  
 1260  
 1265  
 1270  
 1275  
 1280  
 1285  
 1290  
 1295  
 1300  
 1305  
 1310  
 1315  
 1320  
 1325  
 1330  
 1335  
 1340  
 1345  
 1350  
 1355  
 1360  
 1365  
 1370  
 1375  
 1380  
 1385  
 1390  
 1395  
 1400  
 1405  
 1410  
 1415  
 1420  
 1425  
 1430  
 1435  
 1440  
 1445  
 1450  
 1455  
 1460  
 1465  
 1470  
 1475  
 1480  
 1485  
 1490  
 1495  
 1500  
 1505  
 1510  
 1515  
 1520  
 1525  
 1530  
 1535  
 1540  
 1545  
 1550  
 1555  
 1560  
 1565  
 1570  
 1575  
 1580  
 1585  
 1590  
 1595  
 1600  
 1605  
 1610  
 1615  
 1620  
 1625  
 1630  
 1635  
 1640  
 1645  
 1650  
 1655  
 1660  
 1665  
 1670  
 1675  
 1680  
 1685  
 1690  
 1695  
 1700  
 1705  
 1710  
 1715  
 1720  
 1725  
 1730  
 1735  
 1740  
 1745  
 1750  
 1755  
 1760  
 1765  
 1770  
 1775  
 1780  
 1785  
 1790  
 1795  
 1800  
 1805  
 1810  
 1815  
 1820  
 1825  
 1830  
 1835  
 1840  
 1845  
 1850  
 1855  
 1860  
 1865  
 1870  
 1875  
 1880  
 1885  
 1890  
 1895  
 1900  
 1905  
 1910  
 1915  
 1920  
 1925  
 1930  
 1935  
 1940  
 1945  
 1950  
 1955  
 1960  
 1965  
 1970  
 1975  
 1980  
 1985  
 1990  
 1995  
 2000  
 2005  
 2010  
 2015  
 2020  
 2025  
 2030  
 2035  
 2040  
 2045  
 2050  
 2055  
 2060  
 2065  
 2070  
 2075  
 2080  
 2085  
 2090  
 2095  
 2100  
 2105  
 2110  
 2115  
 2120  
 2125  
 2130  
 2135  
 2140  
 2145  
 2150  
 2155  
 2160  
 2165  
 2170  
 2175  
 2180  
 2185  
 2190  
 2195  
 2200  
 2205  
 2210  
 2215  
 2220  
 2225  
 2230  
 2235  
 2240  
 2245  
 2250  
 2255  
 2260  
 2265  
 2270  
 2275  
 2280  
 2285  
 2290  
 2295  
 2300  
 2305  
 2310  
 2315  
 2320  
 2325  
 2330  
 2335  
 2340  
 2345  
 2350  
 2355  
 2360  
 2365  
 2370  
 2375  
 2380  
 2385  
 2390  
 2395  
 2400  
 2405  
 2410  
 2415  
 2420  
 2425  
 2430  
 2435  
 2440  
 2445  
 2450  
 2455  
 2460  
 2465  
 2470  
 2475  
 2480  
 2485  
 2490  
 2495  
 2500  
 2505  
 2510  
 2515  
 2520  
 2525  
 2530  
 2535  
 2540  
 2545  
 2550  
 2555  
 2560  
 2565  
 2570  
 2575  
 2580  
 2585  
 2590  
 2595  
 2600  
 2605  
 2610  
 2615  
 2620  
 2625  
 2630  
 2635  
 2640  
 2645  
 2650  
 2655  
 2660  
 2665  
 2670  
 2675  
 2680  
 2685  
 2690  
 2695  
 2700  
 2705  
 2710  
 2715  
 2720  
 2725  
 2730  
 2735  
 2740  
 2745  
 2750  
 2755  
 2760  
 2765  
 2770  
 2775  
 2780  
 2785  
 2790  
 2795  
 2800  
 2805  
 2810  
 2815  
 2820  
 2825  
 2830  
 2835  
 2840  
 2845  
 2850  
 2855  
 2860  
 2865  
 2870  
 2875  
 2880  
 2885  
 2890  
 2895  
 2900  
 2905  
 2910  
 2915  
 2920  
 2925  
 2930  
 2935  
 2940  
 2945  
 2950  
 2955  
 2960  
 2965  
 2970  
 2975  
 2980  
 2985  
 2990  
 2995  
 3000  
 3005  
 3010  
 3015  
 3020  
 3025  
 3030  
 3035  
 3040  
 3045  
 3050  
 3055  
 3060  
 3065  
 3070  
 3075  
 3080  
 3085  
 3090  
 3095  
 3100  
 3105  
 3110  
 3115  
 3120  
 3125  
 3130  
 3135  
 3140  
 3145  
 3150  
 3155  
 3160  
 3165  
 3170  
 3175  
 3180  
 3185  
 3190  
 3195  
 3200  
 3205  
 3210  
 3215  
 3220  
 3225  
 3230  
 3235  
 3240  
 3245  
 3250  
 3255  
 3260  
 3265  
 3270  
 3275  
 3280  
 3285  
 3290  
 3295  
 3300  
 3305  
 3310  
 3315  
 3320  
 3325  
 3330  
 3335  
 3340  
 3345  
 3350  
 3355  
 3360  
 3365  
 3370  
 3375  
 3380  
 3385  
 3390  
 3395  
 3400  
 3405  
 3410  
 3415  
 3420  
 3425  
 3430  
 3435  
 3440  
 3445  
 3450  
 3455  
 3460  
 3465  
 3470  
 3475  
 3480  
 3485  
 3490  
 3495  
 3500  
 3505  
 3510  
 3515  
 3520  
 3525  
 3530  
 3535  
 3540  
 3545  
 3550  
 3555  
 3560  
 3565  
 3570  
 3575  
 3580  
 3585  
 3590  
 3595  
 3600  
 3605  
 3610  
 3615  
 3620  
 3625  
 3630  
 3635  
 3640  
 3645  
 3650  
 3655  
 3660  
 3665  
 3670  
 3675  
 3680  
 3685  
 3690  
 3695  
 3700  
 3705  
 3710  
 3715  
 3720  
 3725  
 3730  
 3735  
 3740  
 3745  
 3750  
 3755  
 3760  
 3765  
 3770  
 3775  
 3780  
 3785  
 3790  
 3795  
 3800  
 3805  
 3810  
 3815  
 3820  
 3825  
 3830  
 3835  
 3840  
 3845  
 3850  
 3855  
 3860  
 3865  
 3870  
 3875  
 3880  
 3885  
 3890  
 3895  
 3900  
 3905  
 3910  
 3915  
 3920  
 3925  
 3930  
 3935  
 3940  
 3945  
 3950  
 3955  
 3960  
 3965  
 3970  
 3975  
 3980  
 3985  
 3990  
 3995  
 4000  
 4005  
 4010  
 4015  
 4020  
 4025  
 4030  
 4035  
 4040  
 4045  
 4050  
 4055  
 4060  
 4065  
 4070  
 4075  
 4080  
 4085  
 4090  
 4095  
 4100  
 4105  
 4110  
 4115  
 4120  
 4125  
 4130  
 4135  
 4140  
 4145  
 4150  
 4155  
 4160  
 4165  
 4170  
 4175  
 4180  
 4185  
 4190  
 4195  
 4200  
 4205  
 4210  
 4215  
 4220  
 4225  
 4230  
 4235  
 4240  
 4245  
 4250  
 4255  
 4260  
 4265  
 4270  
 4275  
 4280  
 4285  
 4290  
 4295  
 4300  
 4305  
 4310  
 4315  
 4320  
 4325  
 4330  
 4335  
 4340  
 4345  
 4350  
 4355  
 4360  
 4365  
 4370  
 4375  
 4380  
 4385  
 4390  
 4395  
 4400  
 4405  
 4410  
 4415  
 4420  
 4425  
 4430  
 4435  
 4440  
 4445  
 4450  
 4455  
 4460  
 4465  
 4470  
 4475  
 4480  
 4485  
 4490  
 4495  
 4500  
 4505  
 4510  
 4515  
 4520  
 4525  
 4530  
 4535  
 4540  
 4545  
 4550  
 4555  
 4560  
 4565  
 4570  
 4575  
 4580  
 4585  
 4590  
 4595  
 4600  
 4605  
 4610  
 4615  
 4620  
 4625  
 4630  
 4635  
 4640  
 4645  
 4650  
 4655  
 4660  
 4665  
 4670  
 4675  
 4680  
 4685  
 4690  
 4695  
 4700  
 4705  
 4710  
 4715  
 4720  
 4725  
 4730  
 4735  
 4740  
 4745  
 4750  
 4755  
 4760  
 4765  
 4770  
 4775  
 4780  
 4785  
 4790  
 4795  
 4800  
 4805  
 4810  
 4815  
 4820  
 4825  
 4830  
 4835  
 4840  
 4845  
 4850  
 4855  
 4860  
 4865  
 4870  
 4875  
 4880  
 4885  
 4890  
 4895  
 4900  
 4905  
 4910  
 4915  
 4920  
 4925  
 4930  
 4935  
 4940  
 4945  
 4950  
 4955  
 4960  
 4965  
 4970  
 4975  
 4980  
 4985  
 4990  
 4995  
 5000  
 5005  
 5010  
 5015  
 5020  
 5025  
 5030  
 5035  
 5040  
 5045  
 5050  
 5055  
 5060  
 5065  
 5070  
 5075  
 5080  
 5085  
 5090  
 5095  
 5100  
 5105  
 5110  
 5115  
 5120  
 5125  
 5130  
 5135  
 5140  
 5145  
 5150  
 5155  
 5160  
 5165  
 5170  
 5175  
 5180  
 5185  
 5190  
 5195  
 5200  
 5205  
 5210  
 5215  
 5220  
 5225  
 5230  
 5235  
 5240  
 5245  
 5250  
 5255  
 5260  
 5265  
 5270  
 5275  
 5280  
 5285  
 5290  
 5295  
 5300  
 5305  
 5310  
 5315  
 5320  
 5325  
 5330  
 5335  
 5340  
 5345  
 5350  
 5355  
 5360  
 5365  
 5370  
 5375  
 5380  
 5385  
 5390  
 5395  
 5400  
 5405  
 5410  
 5415  
 5420  
 5425  
 5430  
 5435  
 5440  
 5445  
 5450  
 5455  
 5460  
 5465  
 5470  
 5475  
 5480  
 5485  
 5490  
 5495  
 5500  
 5505  
 5510  
 5515  
 5520  
 5525  
 5530  
 5535  
 5540  
 5545  
 5550  
 5555  
 5560  
 5565  
 5570  
 5575  
 5580  
 5585  
 5590  
 5595  
 5600  
 5605  
 5610  
 5615  
 5620  
 5625  
 5630  
 5635  
 5640  
 5645  
 5650  
 5655  
 5660  
 5665  
 5670  
 5675  
 5680  
 5685  
 5690  
 5695  
 5700  
 5705  
 5710  
 5715  
 5720  
 5725  
 5730  
 5735  
 5740  
 5745  
 5750  
 5755  
 5760  
 5765  
 5770  
 5775  
 5780  
 5785  
 5790  
 5795  
 5800  
 5805  
 5810  
 5815  
 5820  
 5825  
 5830  
 5835  
 5840  
 5845  
 5850  
 5855  
 5860  
 5865  
 5870  
 5875  
 5880  
 5885  
 5890  
 5895  
 5900  
 5905  
 5910  
 5915  
 5920  
 5925  
 5930  
 5935  
 5940  
 5945  
 5950  
 5955  
 5960  
 5965  
 5970  
 5975  
 5980  
 5985  
 5990  
 5995  
 6000  
 6005  
 6010  
 6015  
 6020  
 6025  
 6030  
 6035  
 6040  
 6045  
 6050  
 6055  
 6060  
 6065  
 6070  
 6075  
 6080  
 6085  
 6090  
 6095  
 6100  
 6105  
 6110  
 6115  
 6120  
 6125  
 6130  
 6135  
 6140  
 6145  
 6150  
 6155  
 6160  
 6165  
 6170  
 6175  
 6180  
 6185  
 6190  
 6195  
 6200  
 6205  
 6210  
 6215  
 6220  
 6225  
 6230  
 6235  
 6240  
 6245  
 6250  
 6255  
 6260  
 6265  
 6270  
 6275  
 6280  
 6285  
 6290  
 6295  
 6300  
 6305  
 6310  
 6315  
 6320  
 6325  
 6330  
 6335  
 6340  
 6345  
 6350  
 6355  
 6360  
 6365  
 6370  
 6375  
 6380  
 6385  
 6390  
 6395  
 6400  
 6405  
 6410  
 6415  
 6420  
 6425  
 6430  
 6435  
 6440  
 6445  
 6450  
 6455  
 6460  
 6465  
 6470  
 6475  
 6480  
 6485  
 6490  
 6495  
 6500  
 6505  
 6510  
 6515  
 6520  
 6525  
 6530  
 6535  
 6540  
 6545  
 6550  
 6555  
 6560  
 6565  
 6570  
 6575  
 6580  
 6585  
 6590  
 6595  
 6600  
 6605  
 6610  
 6615  
 6620  
 6625  
 6630  
 6635  
 6640  
 6645  
 6650  
 6655  
 6660  
 6665  
 6670  
 6675  
 6680  
 6685  
 6690  
 6695  
 6700  
 6705  
 6710  
 6715  
 6720  
 6725  
 6730  
 6735  
 6740  
 6745  
 6750  
 6755  
 6760  
 6765  
 6770  
 6775  
 6780  
 6785  
 6790  
 6795  
 6800  
 6805  
 6810  
 6815  
 6820  
 6825  
 6830  
 6835  
 6840  
 6845  
 685

<210> 37  
 <211> 65  
 <212> PRT  
 <213> Bacterium  
 5  
 <220>  
 <223> protein RelB-Af1  
 <400> 37  
 10 Met Asn Glu Ala Leu Leu Arg Glu Ile Tyr Ser Glu Val Lys Lys Ile  
     1                    5                    10                    15  
 Arg Glu Lys Ile Glu Gln Leu Glu Glu Leu Ile Ile Pro Ala Glu Lys  
                     20                    25                    30  
 Val Ser Glu Glu Glu Leu Leu Glu Ile Arg Lys Leu Lys Glu Glu Ser  
 15                    35                    40                    45  
 Leu Lys Gly Glu His Val Asp Trp Asp Glu Leu Lys Arg Glu Leu Gly  
             50                    55                    60  
 Val  
 65  
 20  
 <210> 38  
 <211> 72  
 <212> PRT  
 <213> Bacterium  
 25  
 <220>  
 <223> protein RelB-Af3  
 <400> 38  
 30 Met Lys Val Leu Leu Asp Ile Ile Glu Asp Ile Glu Asn Phe Ile Arg  
     1                    5                    10                    15  
 Gln Leu Glu Lys Arg Arg Gly Glu Leu Glu Glu Leu Lys Asp Glu Ile  
                     20                    25                    30  
 Leu Ile Phe Ser Asp Ala Glu Phe Ile Asp Ser Ile Gln Arg Gly Leu  
 35                    35                    40                    45  
 Ser Asp Leu Glu Gln Gly Arg Ser Lys Val Cys Ser Asn Leu Glu Glu  
             50                    55                    60  
 Val Lys Lys Leu Phe Glu Asp Ile  
 65                    70  
 40  
 <210> 39  
 <211> 62  
 <212> PRT  
 <213> Bacterium  
 45  
 <220>  
 <223> protein RelB-Af2  
 <400> 39  
 50 Met Glu Val Ile Gln Ile Ser Lys Asp Glu Leu Glu Glu Ile Ile Glu  
     1                    5                    10                    15  
 Arg Lys Phe Lys Glu Val Leu Ile Lys Ala Leu Met Glu Ile Thr Pro  
                     20                    25                    30  
 Tyr Val Ser Asp Glu Glu Gln Glu Glu Ile Asp Lys Ile Ala Gly Lys  
 55                    35                    40                    45  
 Pro Asp Glu Tyr Glu Gly Glu Phe Glu Glu Trp His Gly Lys  
             50                    55                    60



<210> 40  
 <211> 57  
 <212> PRT  
 <213> Bacterium  
 5  
 <220>  
 <223> protein RelB-Af4  
 <400> 40  
 10 Met Asp Ile Gln Val Ile Lys Gln Ala Val Arg Glu Val Leu Arg Glu  
     1                    5                    10                    15  
 Glu Leu Pro Ser Ile Leu Lys Glu Val Ile Leu Ser Thr Ile Pro Pro  
                     20                    25                    30  
 Asp Glu Pro Glu Ala Asp Glu Lys Gln Phe Val Asp Glu Glu Ile Asn  
 15                    35                    40                    45  
 Glu Asp Asp Tyr Val Lys Phe Asp Glu  
     50                    55  
 <210> 41  
 20 <211> 95  
 <212> PRT  
 <213> Bacterium  
 <220>  
 25 <223> protein RelB-Hp  
 <400> 41  
 Met Pro Asn Thr Thr Asn Lys Asp Tyr Thr Lys Tyr Ser Gln Arg Gln  
     1                    5                    10                    15  
 30 Leu Phe Ser Phe Leu Asn Ser Ile Lys Thr Lys Gln Lys Arg Ala Leu  
                     20                    25                    30  
 Glu Lys Leu Lys Glu Ile Gln Ala Gln Lys Gln Arg Ile Lys Lys Ala  
                     35                    40                    45  
 Leu Gln Phe Lys Ala Leu Asn Leu Thr Glu Asn Gly Tyr Thr Ile Glu  
 35                    50                    55                    60  
 Glu Glu Arg Glu Ile Leu Ala Arg Ala Lys Asp Thr Lys Asn Arg Leu  
     65                    70                    75                    80  
 Cys Phe Lys Ser Ile Glu Asp Phe Lys Lys His Cys Glu Asn Leu  
                     85                    90                    95  
 40  
 <210> 42  
 <211> 86  
 <212> PRT  
 <213> Bacterium  
 45  
 <220>  
 <223> protein RelB-syneco  
 <400> 42  
 50 Met Met Arg Ala Phe Glu Val Met Ala Thr Val Lys Asp Ser Lys Gln  
     1                    5                    10                    15  
 Leu Leu Leu Asp Ser Asp Leu His Trp Asn Thr Ser Arg Val Lys Val  
                     20                    25                    30  
 Ile Ile Leu Glu Ser Asp Glu Leu Ala Ser Lys Gly Ser Glu Phe Asp  
 55                    35                    40                    45  
 Pro Asp Asp Thr Pro Val Glu Glu Ile Lys Val Ser Leu Arg Lys Ala  
     50                    55                    60  
 Leu Glu Glu Tyr Lys Gln Gly Lys Arg Ile Pro Val Glu Asn Met Trp  
     65                    70                    75                    80

Glu Gly Ile Asp Val Glu  
85

5 <210> 43  
<211> 85  
<212> PRT  
<213> Bacterium

10 <220>  
<223> protein RelB-BT

<400> 43  
Met Ala Ile Arg Lys Asp Glu Leu Tyr Arg Leu Ile Asp His Leu Asp  
1 5 10 15  
15 Gln Gln Asp Glu Lys Ala Ala Phe Asp Phe Leu Glu Phe Leu Val Gln  
20 25 30  
Arg Ser Arg Arg Lys Pro Lys Glu Trp Glu Lys Ile Asp Met Ala Asp  
35 40 45  
20 Pro Asp His Glu Pro Leu Ser Thr Gln Glu Leu Glu Gln Leu Asn Ser  
50 55 60  
Glu Glu Gly Tyr Val Ser Gly Glu Asp Ala Lys Arg Glu Phe Gly Leu  
65 70 75 80  
Gln Ile Asp Leu Pro  
85

25 <210> 44  
<211> 1280  
<212> DNA  
<213> E. coli plasmid P307

30 <400> 44  
gagtatcata ttaggatacg ggtgggtgac gccacacctt gccatagaac ggacattcat 60  
tgatgccatg ccagaatgga cgttcaggtt attccgtcca gttctgctgg caacgcgaga 120  
tctccctgga tatagtgatg ccacagcaaa gcgctcaaac agggataata tgatggaaat 180  
35 caaggetcaa cagttttgtc acatcaacgg ggcggaagt ccttactgac aacggacaac 240  
aaggtatggg cggcgtggcg ggtatcggtt ccacgactga aaagcatcag gggcgctggg 300  
cggaagcgat ttttgcaaac tgcgcggaac tggataacga ccagcttaac gagatcatcg 360  
agtgggttcg gctctatcag cgctgaatgc cactatcagg ctgcgcaagc ggcctttttt 420  
acgccccttg ttttaattccc gcactacctg gacgttcagg tgattctgtc catctgtaca 480  
40 aaaaacaata aaagacttgt rbmrnataac aggtcatgta aggagtatct ttgagactgg 540  
ttaaacagtc ttgaaasdrt arttrbgttg cctatgccta acattattct cagtgtatca 600  
agcgccagtg tcagcgagct gaagaaaaac ccgatggcga cagtcagcgc cggatgatgg 660  
ttcccggtcg ctatcctgaa ccgtaatcag cctgctttct actgtgtacc cgcagagctg 720  
tacgaaaaga tgcttgatgc cctagacgat caggagtggg ttaaasdctg gtagccgaac 780  
45 gcagcaacca accgctgcat gatgtagatc tggataandr bstartrata cctatgaggt 840  
atcaggtaaa attcagggaa gatgcgctga aagagtggca aaaactggac aaggctattc 900  
agcaacagtt tgcgaaaaag ctaaaaaagt gctgtgacaa tccgcatatt ccttccgcaa 960  
aactgcgtgg gataaaggac tgctacaaaa taaaattacg tgcgtcaggt ttctgcctgg 1020  
tctatcaggt gattgacgaa caattaatta tcgctgttgt agctgtgggt aaacgtgndr 1080  
50 agcgagtgta cgttttataat cttgccagcg aaagaatgag ataaaagcaa taaacacaga 1140  
aagtactctt ggcgttatgg ggtaatgcaa agtatgagtc gttaggggaa ttgcctggat 1200  
aattcgccga tggaaagagt ctttcgcagc cttaaaagtg aatggcttcc gaaaggtggg 1260  
tatgggtgatt ttagccatgc 1280

55 <210> 45  
<211> 1168  
<212> DNA  
<213> B. thuringiensis

	<400> 45	
	ctcgtttttt ctggttggtac aaacttaatt gattttgaat aatttggttg taccagtcct	60
	ttttgcttag cccagtc aaa ataacgtttg attgaattaa tgcgccggtt aatcgtagaa	120
	ggttttagta atcttgtaac ttgcatatgc ctcgatatc gagcaatagt gcgagcggta	180
5	acttctattg gatgaaaaag agtatcctca gcatgttttc cccacacatt ttcaaacc aa	240
	aatacaaaaat cttttaaatc actcgtatat tcttttagtg tttttgtatg caaatctcct	300
	tcttgagata agctagaaat aaaatcggaa atcaaagatg ttgcttgatg agaaattggt	360
	ttagtggaat gcataaatac ctctcttttt attgacttac rbbtmrnaat tagcggacat	420
	gataattttaa tcttatcaat tatgttagcg gacatcaaac atttattttc ccacacttca	480
10	tgtccactaa tattaattag tggacattr sdstartrbta aaactatctc gaaagtaggt	540
	gtaacacatg gctattcgt aagatgaatt gtatcgggtta attgatcacc tggatcaaca	600
	agatgaaaaa gcagcatttg actttttaga atttcttggt caacgggtcaa gaagaaaacc	660
	taaagaatgg gaaaaaattg atatggcaga tcttgatcat gaaccgctgt ctacacaaga	720
	gttagaacag ttaaacagtg aagaaggata tgtatcaggg gaggacgcnd rbstartraa	780
15	aacgtgaatt cggtacataa attgatttac cataagtcgg cggtgaaatt tattgcaaag	840
	caagaaaaag ggattcaaaa aagaattgca gaaggattga agggacttct taagattcct	900
	cctgaaggag atattaaaaag tatgaaagggt tacacagaac tatatcgatt acggattgga	960
	acctttcgaa ttttatttga aataaatcat gatgagaaag tcatatacat acaagcaatt	1020
	ggaaatcgnd rtggtgacat ctataaataa ggcaaacatg catttttaaa agaaagggtct	1080
20	tctgaatcga agaaccttcc tttttgtgt gcgaataatg tccgctaattg cttgttgctg	1140
	gattctgttc cattgctaca catacccc	1168

	<210> 46
	<211> 1024
25	<212> DNA
	<213> Methanococcus jannaschii

	<400> 46	
	ccgataccgt tgctggagac atagctggag ctttgaaggc ggagaagctt attttaataa	60
30	cagatgttga tgggaataatg gatgatataa ataatccaga gacgttgcat agaaaaataa	120
	cagcttcaga actaaaagaa atgatagaag atggaagaat aaaggagggg atgattccaa	180
	aggctgaaa tgctttatat gccttagagc atggagttaa gagcgttcat ataataaatg	240
	gaaagattcc tcatgctttg ttggttgaga tatttacaga ggagggtatt gggacgatga	300
	taacaagaga ttaaagtttt tatattataa actacttaag aattaaaata startrbmag	360
35	acaaataagg ggataactat gctcaatata aacaaagaga tagcacaat agaaactgaa	420
	ttgaatgaat tgaaaaaatt gagagatgaa atctctgaaa ggattgaaaa attagaaata	480
	aagttattaa aattgaaagc attagctatt ccagaggagg aatttgaaga ggattatgaa	540
	gaaattatag aagatgttaa aaaatctctg gataaaaaag agactgtgcc agcagaagag	600
	gctttgaand rbmstartrm agaattggga ttattatgaa gtttaacgtt gagatacata	660
40	aaagagtctt aaaagattta aaggatttgc ctccctcaaa cttaaagaag tttaaagaac	720
	taatagaac attaaaaacc aatcccattc caaaagaaaa atttgatatt aaaagattaa	780
	aaggcagtga tgaggtttat agagtttaga ttggaaaatt tagagttcaa tatgttgtt	840
	tatgggatga tagaataata ataattagaa ndrmagataa gtagaagaga agggagcttat	900
	aaaaatccct aagctattaa aaattcta at ggctacattt ttatatctct tttcttaatt	960
45	caaatagaaa aaacagattc ggctgatacc atgattattc ttttagattt aaatggaaca	1020
	atag	1024

	<210> 47
	<211> 28
50	<212> DNA
	<213> Artificial Sequence

	<220>
55	<223> Primer MJ-relE/2CW was used for the amplification of the relE gene of M. jannaschii from genomic DNA

<400> 47  
ccccgaatt cgcatgcgcc attagaat

- 5       <210> 48  
      <211> 37  
      <212> DNA  
      <213> Artificial Sequence
- <220>  
      <223> Primer MJ-reE/1CW was used for the amplification  
            of the relE gene of M. jannaschii from genomic DNA
- 10       <400> 48  
      cccccgatc cgagctcgag gctttgaaag aattggg 37
- 15       <210> 49  
      <211> 38  
      <212> DNA  
      <213> Artificial Sequence
- 20       <220>  
      <223> Primer relB-M.jannCW was used for the  
            PCR-amplification of relB and relE from M.  
            jannashii
- 25       <400> 49  
      ccccgatcc gtcgacgaca aataagggga taactatg 38
- 30       <210> 50  
      <211> 32  
      <212> DNA  
      <213> Artificial Sequence
- 35       <220>  
      <223> Primer relE-Sp2/cw was used for the  
            PCR-amplification of the gene relESP2 from genomic  
            DNA of S. pneumoniae
- 40       <400> 50  
      ccccgatcc gatgcatgat ttaggcttga ag 32
- 45       <210> 51  
      <211> 34  
      <212> DNA  
      <213> Artificial Sequence
- 50       <220>  
      <223> Primer relE-Sp2/ccw was used for the  
            PCR-amplification of the gene relESP2 from genomic  
            DNA of S. pneumoniae
- 55       <400> 51  
      ccccgaattc gaatgaaaat ttacttgaaa aaag 34
- <210> 52  
      <211> 58  
      <212> DNA  
      <213> Artificial Sequence
- <220>  
      <223> Primer relEk12 was used for the PCR-amplification  
            of DNA-fragments comprising genes relEK-12,

## releP307 and releMj

5                   <400> 52  
tgtaatacga ctcactatag ataaggagtt ttataaatgg cgtattttct ggattttg                   58

                  <210> 53  
                  <211> 19  
                  <212> DNA  
                  <213> Artificial Sequence

10                   <220>  
                  <223> Primer P2 was used for the PCR-amplification of  
                          DNA-fragments comprising genes releK-12, releP307  
                          and releMj

15                   <400> 53  
caccttcggt gcgaaacag                   19

20                   <210> 54  
                  <211> 58  
                  <212> DNA  
                  <213> Artificial Sequence

25                   <220>  
                  <223> Primer releP307 was used for the PCR-amplification  
                          of DNA-fragments comprising genes releK-12,  
                          releP307 and releMj

30                   <400> 54  
tgtaatacga ctcactatag ataaggagtt ttataaatga ggtatcaggt aaaattca                   58

                  <210> 55  
                  <211> 20  
                  <212> DNA  
35                   <213> Artificial sequence

                  <220>  
                  <223> Primer P4 was used for the PCR-amplification of  
                          DNA-fragments comprising genes releK-12, releP307  
                          and releMj

40                   <400> 55  
ctttccatcg gcgaattatc                   20

45                   <210> 56  
                  <211> 58  
                  <212> DNA  
                  <213> Artificial Sequence

50                   <220>  
                  <223> Primer releMj was used for the PCR-amplification  
                          of DNA-fragments comprising genes releK-12,  
                          releP307 and releMj

55                   <400> 56  
tgtaatacga ctcactatag ataaggagtt ttataaatga agtttaacgt tgagatac                   58


	<210> 57	
	<211> 20	
	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
	<223> Primer P6 was used for the PCR-amplification of	
	DNA-fragments comprising genes relEK-12, relEP307	
	and relEMj	
10	<400> 57	
	atcatggtat cagccgaatc	20
	<210> 58	
15	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
20	<223> Primer S-RelE was used for the PCR-amplification	
	of the RelE coding region from the plasmid pMG223	
	<400> 58	
	taggtacccat ggcgtatttt ctgg	24
25	<210> 59	
	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Primer AS-RelE was used for the PCR-amplification	
	of the RelE coding region from the plasmid pMG223	
35	<400> 59	
	gagaccccccac actaccatcg gcg	23

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A. The indications made below relate to the microorganism referred to in the description</b> on page <u>33</u> , line <u>13</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input checked="" type="checkbox"/></span>	
Name of depositary institution DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH	
Address of depositary institution (including postal code and country) Mascheroder Weg 1B D-38124 Braunschweig Germany	
Date of deposit 30 April 1998	Accession Number DSM 12157
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
As regards the respective Patent Offices of the respective designated states, the applicants request that a sample of the deposited microorganisms only be made available to an expert nominated by the requester until the date on which the patent is granted or the date on which the application has been refused or withdrawn or is deemed to be withdrawn.	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input type="checkbox"/> This sheet was received with the international application	
Authorized officer	

For International Bureau use only	
<input checked="" type="checkbox"/> This sheet was received by the International Bureau on:	
<b>13 SEP 1999</b>	
Authorized officer 	

INDICATIONS RELATING TO DEPOSITED MICROORGANISMS  
(PCT Rule 12bis)

5

Additional sheet

In addition to the microorganism indicated on page 33 of the description, the following microorganisms have been deposited with

- 10 DSM-Deutsche Sammlung von Mikroorganismen und Cellkulturen GmbH  
Mascheroder Weg 1b, D-38124 Braunschweig, Germany

on the dates and under the accession numbers as stated below:

15

Accession number	Date of deposit	Description Page No.	Description Line No.
20 DSM 12158	30 April 1998	33	32
DSM 12159	30 April 1998	35	10
DSM 12160	30 April 1998	34	31
DSM 12161	30 April 1998	33	21

25

For all of the above-identified deposited microorganisms, the following additional indications apply:

30

As regards the respective Patent Offices of the respective designated states, the applicants request that a sample of the deposited microorganisms stated above only be made available to an expert nominated by the requester until the date on which the patent is granted or the date on which the application has been refused or withdrawn or is deemed

35 to be withdrawn.